

THE ROLE OF BDNF IN 17 β -ESTRADIOL-INDUCED FACILITATION OF EXTINCTION
OF COCAINE SEEKING

by

Hanna Yousuf

A Thesis Submitted in
Partial Fulfillment of the
Requirements for the Degree of

Master of Science

in Psychology

at

The University of Wisconsin-Milwaukee

December 2016

ABSTRACT

THE ROLE OF BDNF IN 17 β -ESTRADIOL-INDUCED FACILITATION OF EXTINCTION OF COCAINE SEEKING

by

Hanna Yousuf

The University of Wisconsin Milwaukee, 2016
Under the Supervision of Professor Devin Mueller

Females are more susceptible to drug dependence than males, and these differences in addictive behaviors are mediated, in large part, by 17 β -estradiol (E₂). E₂ enhances memory in a variety of behavioral paradigms and may act to enhance the formation of drug-related memories. Interestingly, due to its mnemonic effects, E₂ promotes drug seeking, but also may aid in extinguishing drug seeking behaviors. Treatment of addiction is modeled through extinction, and the infralimbic medial prefrontal cortex (IL-mPFC) is responsible for inhibiting drug seeking after extinction. However, the cellular mechanisms by which E₂ facilitates extinction remain unknown. Previous work suggests that the memory-enhancing effects of E₂ are mediated by brain-derived neurotrophic factor (BDNF) and its receptor, tropomyosin-related kinase B (TrkB). Thus, we hypothesized that E₂-facilitated extinction is dependent upon BDNF/TrkB signaling. To identify this mechanism, we used the conditioned place preference paradigm (CPP) in which rats are trained to associate one chamber, but not another, with cocaine. Following conditioning, ovariectomized rats were systemically administered E₂ prior to each extinction trial. Furthermore, IL-mPFC infusions of a Trk receptor antagonist did not impair, but facilitated

extinction. Systemic injections of a TrkB receptor antagonist and E₂ did impair extinction, whereas injections of vehicle and E₂ facilitated extinction. The results of this experiment are in agreement with our hypothesis, however, an E₂-facilitated extinction was not consistently observed in this study. In order to determine the cellular mechanism by which E₂ increases intrinsic excitability in IL-mPFC pyramidal neurons, we performed patch-clamp electrophysiology while bath-applying Trk receptor antagonists. Our results revealed that blockade of Trk receptors attenuated E₂-induced increase in intrinsic excitability. These results indicate that E₂ potentiates intrinsic excitability via BDNF/TrkB signaling in IL-mPFC pyramidal neurons. This study demonstrates that optimizing E₂ levels and/or augmenting BDNF function could enhance therapeutic interventions in alleviating learning and memory disorders such as drug addiction.

TABLE OF CONTENTS

ABSTRACT	ii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vii
LIST OF TABLES	viii
ACKNOWLEDGEMENTS	ix
INTRODUCTION	1
Sex differences in addiction	1
Extinction of drug seeking	3
Effects of 17 β -estradiol (E ₂) on learning memory	5
<i>Effects of E₂ on hippocampal memory</i>	6
<i>Effects of E₂ on extinction memory</i>	8
<i>Effects of E₂ on intrinsic excitability</i>	11
Mechanisms underlying the role of E ₂ in learning and memory	12
<i>Interactions between E₂ and BDNF</i>	13
Clinical relevance	16
AIMS	17
Aim 1	17
Aim 2	17
METHODS	19
Subjects and Surgery	19
Drugs.....	20
Conditioned place preference.....	20

<i>Place preference apparatus</i>	20
<i>Conditioning and Testing</i>	21
<i>Experiment one</i>	22
<i>Experiment two</i>	22
<i>Experiment three</i>	23
<i>Histology</i>	23
<i>Data analysis</i>	23
Patch-clamp electrophysiology.....	24
<i>Experimental design</i>	24
<i>Immunohistochemistry</i>	25
<i>Electrophysiological data analysis</i>	26
RESULTS	27
Trk receptor blockade in IL-mPFC facilitated extinction of a cocaine CPP.....	27
Trk receptor blockade in IL-mPFC did not impair extinction of a cocaine CPP in the absence of E ₂	31
E ₂ -induced facilitation of extinction of a cocaine CPP was disrupted by systemic injections of TrkB receptor antagonist.....	33
E ₂ potentiates IL-mPFC pyramidal neuron excitability.....	35
Trk receptor blockade prevented E ₂ -induced potentiation of IL-mPFC pyramidal neuron excitability	39
E ₂ does not facilitate consolidation of extinction of a cocaine CPP.....	41
DISCUSSION	42

Role of Trk receptors in extinction of cocaine seeking.....	43
E ₂ potentiates intrinsic excitability in IL-mPFC pyramidal neurons via a BDNF- dependent mechanism.....	47
Future directions.....	50
CONCLUSION.....	51
REFERENCES.....	52

LIST OF FIGURES

Figure 1, Page 29: A single IL-mPFC infusion of non-selective Trk receptor antagonist facilitated extinction of a cocaine CPP

Figure 2, Page 30: Multiple IL-mPFC infusions of non-selective Trk receptor antagonist facilitated extinction of a cocaine CPP

Figure 3, Page 32: A single IL-mPFC infusion of non-selective Trk receptor antagonist did not disrupt extinction of a cocaine CPP in the absence of E₂

Figure 4, Page 34: Injections of selective TrkB receptor antagonist disrupted E₂-induced facilitation of extinction of a cocaine CPP

Figure 5, Page 36: Bath-application of E₂ potentiates IL-mPFC pyramidal neuron excitability in slices from OVX rats that were primed with E₂

Figure 6, Page 38: Bath-application of E₂ potentiates IL-mPFC pyramidal neuron excitability in slices from OVX rats that were not primed with E₂

Figure 7, Page 40: Blockade of Trk receptors prevent E₂-induced potentiation of IL-mPFC neuronal intrinsic excitability

Figure 8, Page 41: Post-injections of E₂ did not facilitate consolidation of extinction of a cocaine CPP

LIST OF TABLES

Table 1, Page 37: Effects of E₂ on intrinsic excitability of IL-mPFC pyramidal neurons in slices from OVX rats that were primed with E₂

Table 2, Page 38: Effects of E₂ on intrinsic excitability of IL-mPFC pyramidal neurons in slices from OVX rats that were not primed with E₂

Table 3, Page 40: Effects of E₂ and K-252a on intrinsic excitability of IL-mPFC pyramidal neurons

ACKNOWLEDGEMENTS

I would like to thank my advisor Dr. Devin Mueller for his guidance and help throughout this project. I would also like to thank my committee members, Dr. Karyn Frick and Dr. James Moyer for their advice and feedback on my thesis. Thanks to Chad Smies for his assistance during the behavioral experiments. I am extremely grateful to Pat Reily for his ingenuity and help with the CPP boxes. I want to extend my sincerest gratitude to Jen Tuscher and Madalyn Hafenbreidel for their advice and friendship. Finally, thank you to Nick Fedorchuk for always supporting me throughout the process.

INTRODUCTION

Sex differences in addiction

In 2014, The National Survey on Drug Use and Health (NSDUH) reported that approximately 1.5 million Americans aged 12 and older currently used cocaine. The prevalence of addiction is reported to be greater in men compared to women (55.1% vs 44% overall, 18.1% vs 11.4% for cocaine respectively, SAMSHA 2010), therefore, most treatment programs and research on drug abuse focus on the male population. However, cocaine dependence is greater in females than males at a younger age (Kandel, Chen, Warner, Kessler, & Grant 1997; Weiss, Martinez-Raga, Griffin, Greenfield, & Hufford, 1997; Chen & Kandel, 2002), women progress more rapidly from casual drug use to dependence (McCance-Katz, Carroll, & Rounsaville, 1999; Westermeyer & Boedicker, 2000), they engage in binge-like patterns of abuse (Becker & Hu, 2008; Brady & Randall, 1999; Lynch, Roth, & Carroll, 2002), and tend to have greater lifetime use compared to men (Lejuez, Bornovalova, Reynold, Daughters, & Curtin, 2007). In terms of relapse, women have shorter periods of abstinence and are more susceptible to relapse following stressful life events or depression (McKay, Rutherford, Cacciola, Kabasakalian-McKay, & Alterman 1996; Elman, Karlsgodt, & Gastfriend, 2001; Back, Brady, Jackson, Salstrom, & Zinzow, 2005). Psychosocial factors may contribute to gender differences in cocaine-related behaviors, however, preclinical studies suggest that biological factors may also play a significant role in sex differences in cocaine seeking.

Similar to humans, adult female rats are more sensitive to cocaine compared to adult male rats. In the self-administration paradigm, rats are trained to lever press for intravenous infusions of commonly abused drugs. Drug reward is usually presented in conjunction with a light or tone, which becomes a conditioned stimulus (CS+), as an association with a drug reinforcement is

developed (Davis & Smith, 1974; Lynch, Nicholson, Dance, Morgan, & Foley, 2010). Importantly, female rats self-administer cocaine for longer periods and have higher levels of cocaine intake than males (Lynch & Taylor, 2004; Larson, Anker, Gliddon, Fons, & Carroll, 2007; Hu & Becker, 2008; Perry, Westenbroek, & Becker, 2013). Sex differences in addiction have also been observed in the cocaine sensitization paradigm. In this paradigm, repeated exposure to cocaine results in enhanced behavioral responsiveness with subsequent drug administration (Robinson & Becker, 1986; Strakowski, Sax, Setters, & Keck, 1996). Female rats tend to exhibit a greater magnitude of sensitization or rotational behaviors compared to male rats (Hu & Becker, 2003). These sex differences in cocaine seeking have been consistent with findings using another well-established paradigm called the conditioned place preference (CPP) paradigm. In this model, repeated pairings of a drug with a context produces a significant preference for the cocaine-paired environment. Preference is measured by time spent in the drug-paired context compared to time spent in the vehicle-paired context. The preference indicates the association of the environment with the rewarding properties of the drug (Tzschentke, 1998; Tzschentke, 2007). At low doses of cocaine, female rats develop faster associations between the environmental context and cocaine (Russo et al., 2003). Overall, in multiple paradigms, female rats appear to be more vulnerable to the effects of cocaine compared to their male counterparts.

The biological basis for sex-specific differences in cocaine addiction are complex. However, considerable preclinical data suggest that they may be attributed to estrogens (Carroll, Lynch, Roth, Morgan, & Cosgrove, 2004; Lynch et al., 2002; Becker & Hu, 2008). A majority of studies regarding the contribution of estrogens in various behavioral paradigms have been conducted in ovariectomized (OVX) female rodents to provide better experimental control over circulating ovarian hormones through experimenter-administered hormones. For example, in the

self-administration paradigm, OVX female rats treated with the potent estrogen, 17 β -estradiol (E₂), facilitated acquisition of cocaine self-administration (Hu, Crombag, Robinson, & Becker, 2004; Roberts, Bennett, & Vickers, 1989; Lynch, Roth, Mickelberg, & Carroll, 2001; Jackson, Robinson, & Becker, 2006). Similarly, OVX rats treated with estradiol benzoate demonstrated significantly higher cocaine-induced locomotor sensitization (Perrotti et al., 2001; Peris, DeCambre, Coleman-Hardee, & Simpkins, 1991; Sircar & Kim, 1999; Hu & Becker, 2003). However, few studies have used the CPP paradigm to investigate the role of estrogens in cocaine-associated learning. Treatment of estradiol benzoate to OVX rats resulted in a significantly higher preference for the cocaine-paired chamber compared to OVX rats that received vehicle treatment (Bobzean, Dennis, & Perrotti, 2014). These data confirm that cocaine has greater abuse liability in females than males, and that E₂ may mediate these effects.

Extinction of drug seeking

One of the major obstacles to overcoming addiction in both sexes is the propensity to relapse. The rewarding properties of drugs become associated with cues, and when these cues are subsequently encountered, they can trigger drug seeking months or years after drug cessation. However, sex differences are observable during the relapse cycle, as women report shorter periods of abstinence between cocaine uses than men (Kosten, Gawin, Kosten, & Rounsaville, 1993; Ignjatova & Raleva, 2008) and are more likely to relapse after stressful or depressive life events compared to men (McKay et al., 1996; Elman et al., 2001; Back et al., 2005). This effect is also observed in female rodents who show greater cocaine-induced reinstatement than male rodents (Lynch & Carroll, 2000; Kippen, 2005; Kerstetter, Aguilar, Parrish, & Kippen, 2008; Anker & Carroll, 2010). One possible approach to treat maladaptive drug-associated memories is through extinction, which is the formation of a new inhibitory memory that results in the

reduction of learned responses to conditioned drug cues (Franken, deHaan, van der Meer, Haffmans, & Hendriks, 1999; Millan, Marchant, & McNally, 2011; Torregrossa & Taylor, 2013). During extinction, an organism learns that a CS, which is the drug-associated cue, no longer predicts the unconditioned stimulus (i.e., the drug; Bouton, Garcia-Gutierrez, Zilski, & Moody, 2006; Eisenberg & Dudai, 2004; Myers, Carlezon, & Davis, 2011). Furthermore, this results in a reduction in response to drug-paired cues and contexts, as well as a decrease in operant responses that were previously reinforced by a drug reward (Chesworth & Corbit, 2015). As mentioned, extinction does not erase memories of conditioned behavior, but rather is new learning which acts to mask or inhibit original learning (Bouton & Todd, 2014). Similar to other forms of memory, the new inhibitory memory is formed in stages. First, the memory is acquired into short-term storage, and then is consolidated into long-term storage. Subsequently, the memory is retrieved, and then is reconsolidated back into long-term storage (Quirk & Mueller, 2008). Therefore, acquisition of extinction can be manipulated prior to the first extinction trial, whereas consolidation of extinction can be manipulated immediately following the initial extinction session (Millan et al., 2011). Similarly, retrieval of an extinction memory can be manipulated before subsequent extinction trials, and reconsolidation can be manipulated immediately after a subsequent extinction trial (Millan et al., 2011).

The neural circuitry for extinction learning includes the infralimbic medial prefrontal cortex (IL-mPFC), dorsal hippocampus (DH) and nucleus accumbens (NAc). In particular, IL-mPFC has been shown to have a crucial role in extinction of drug seeking. In the self-administration paradigm, local infusions of inactivating agents in the IL-mPFC such as gamma-Aminobutyric acid (GABA) receptor agonists reinstated the previously extinguished drug seeking response (Peters, LaLumiere, & Kalivas, 2008). IL-mPFC also plays an important role in the

consolidation of the extinction memory. Inactivation of IL-mPFC via GABA receptor agonists following daily extinction sessions impaired extinction retention (LaLumiere, Niehoff, & Kalivas, 2010). Moreover, potentiating AMPA receptors in IL-mPFC after extinction training augmented extinction retention in a model of cocaine seeking (LaLumiere et al., 2010). Taken together, these findings highlight the role of IL-mPFC in regulating the consolidation and expression of extinction of drug seeking.

Effects of 17 β -estradiol (E₂) on learning and memory

As suggested earlier, extinction does not eliminate or result in forgetting of the initial conditioned response; rather, extinction involves new learning that the CS no longer predicts the reinforcer (Bouton et al., 2006; Milad & Quirk, 2002; Santini, Muller, & Quirk, 2001). A wealth of scientific literature has shown that E₂ can enhance learning and memory through the activation of signaling cascades and modulation of synapse structure and function, which contributes to the rewiring of neural circuits (Woolley, 2007; Srivastava et al., 2011; Frick, 2015). E₂ also enhances drug-associated memories in rodents and women (Lynch et al., 2001; Larson et al., 2007; Segarra et al., 2014). Therefore, since extinction is a form of learning, E₂ may have the potential to enhance extinction of cocaine seeking.

Estrogens can mediate their effects via two different modes of action: the classical “genomic” mechanism, and the rapid “non-genomic” cell-signaling mechanism. In the genomic mode of action, estrogens bind to intracellular estrogen receptors (ERs), such as ER α and ER β , which act as nuclear transcription factors. The binding of estrogens to ERs causes estrogen-receptor complexes to bind to estrogen response elements (ERE) in the DNA to influence gene transcription. In the non-genomic cell signaling mode of action, estrogens can bind and rapidly activate ERs (e.g. ER α , ER β , G-coupled; Srivastava & Evans, 2013) or neurotransmitter receptors

(e.g. metabotropic glutamate receptors or N-methyl-D-aspartate receptors; Lewis, Kerr, Orr, & Frick, 2008; Boulware, Heisler, & Frick, 2013) located at the cell membrane, which then go on to activate cell-signaling pathways that eventually lead to changes in gene expression and epigenetic alterations (Levin, 2005; Vasudevan & Pfaff, 2007; Frick, Kim, Tuscher, & Fortress, 2015). ERs are expressed in many brain regions, including the hypothalamus, entorhinal cortex, perirhinal cortex, hippocampus, amygdala, prefrontal cortex, and cerebellum (Shughrue, Scrimo, & Merchenthaler, 1998; Kuiper, Shughrue, Mechenthaler, & Gustafsson, 1998; Osterlund, Gustafsson, Keller, & Hurd, 2000; Shughrue & Merchenthaler, 2001; Spencer et al., 2008) and both receptors are located in regions that are critical for the neural circuitry of extinction. Activation of these two types of ERs via E₂ may have differential behavioral consequences such as initiating copulatory behaviors, and improving performance in spatial learning and memory tasks (Luine, Richards, Wu, & Beck, 1998; Brann, Dhandapani, Wakade, Mahesh, & Khan, 2007; Clipperton-Allen, Almey, Melicherick, Allen, & Choleris, 2011; Cornil & Charlier, 2010; Almey et al., 2014).

Effects of E₂ on hippocampal memory

The effects of E₂ on cognitive function have been extensively examined in rodent models and a majority of research has focused on how E₂ regulates hippocampal plasticity and memory. However, there are limited studies that examine the role of E₂ in IL-mPFC and extinction. DH is involved in the circuit that mediates extinction of drug seeking (Peters, Kalivas, & Quirk, 2009). Furthermore, DH has been implicated in extinction as temporary inactivation of this region impairs extinction of conditioned fear (Corcoran & Maren, 2001). Therefore, literature on the effects of E₂ in the hippocampus can give us insight into how E₂ might act in the IL-mPFC to facilitate extinction of cocaine seeking. For example, OVX rodents that have a dramatic loss of

circulating E₂ show impaired performance on hippocampal-dependent tasks such as Morris water maze (Gresack & Frick, 2006; Markham, Pych, & Juraska, 2002; Lowry, Pardon, Yates, & Juraska, 2010; Daniel, Roberts, & Dohanich, 1999; Monteiro, de Mattos, Ben, Netto, & Wyse, 2008), radial arm maze (Fader, Johnson, & Dohanich, 1999; Daniel, Fader, Spencer, & Dohanich, 1997; Bimonte & Denenberg, 1999; Heikkinen, Puolivali, Liu, Rissanen, & Tanila, 2002; Gibbs & Johnson, 2008), object placement, and object recognition (Luine, Jacome, & MacLusky, 2003; Wallace, Luine, Arellanos, & Frankfurt, 2006; Hamilton et al., 2011; Fonseca et al., 2013).

Exogenous treatment of E₂ to young OVX rodents has generally been shown to improve spatial memory. For example, acute or chronic systemic administration of E₂ enhances spatial memory in the Morris water maze, radial arm maze, and object placement (O'Neal, Means, Poole, & Hamm, 1996; Daniel et al., 1997; Fader et al., 1999; Luine et al., 1998; Bimonte & Denenberg, 1999; Sandstrom & Williams, 2001; Luine et al., 2003; Li et al., 2004; Gresack & Frick, 2004; Walf, Rhodes, & Frye, 2006; Inagaki, Gautreaux, & Luine, 2010; Tuscher, Fortress, Kim, & Frick, 2015). Exogenous E₂ can also enhance object recognition memory (Luine et al., 2003; Gresack & Frick, 2006; Lewis et al., 2008; Jacome et al., 2010; Frick, Fernandez, & Harburger, 2010; Tuscher et al., 2015). To validate the role of the hippocampus in spatial memory and object memory consolidation, intrahippocampal infusions of E₂ enhance both forms of memory (Packard, Kohlmaier, & Alexander, 1996; Packard & Teather, 1997; Fernandez et al., 2008; Pereira, Bastos, de Souza, Ribeiro, & Pereira, 2014; Zhao, Fan, & Frick, 2010; Fortress, Fan, Orr, Zhao, & Frick, 2013; Boulware et al., 2013). A similar pattern has been observed when ER-specific agonists were administered to rodents. Administration of a ER β selective agonist, diarylpropionitrile (DPN), but not ER α agonist, propyl pyrazole triol (PPT), mimicked the mnemonic effects of E₂ in the radial arm maze (Liu et al., 2008), Morris water maze (Rhodes &

Frye, 2006), object placement, and object recognition (Jacome et al., 2010; Walf et al., 2006, Walf, Koonce, & Frye, 2008; Frick et al., 2010) in OVX rodents. Conversely, studies have also reported that E₂ either does not improve or, in some cases, inhibits performance on some memory tasks. This variability may be due to dose, duration of treatment, age at the onset of treatment, route of administration, and difficulty of task (Dohanich, 2002; Frick, 2009; Acosta et al., 2009; Holmes, Wide, & Galea, 2002; Gresack & Frick, 2006; Markham & Juraska, 2007; Markowska & Savonenko, 2002; Garza-Meilandt, Cantu, & Claiborne, 2006). Taken together, E₂ has been shown to improve memory in a variety of behavioral tasks dependent on the hippocampus and may act as a key mediator underlying extinction of cocaine seeking in females.

Effects of E₂ on extinction memory

E₂ modulates neural plasticity which can lead to strengthened learning and memory. Furthermore, E₂ can influence neural plasticity within the circuitry underlying extinction of fear conditioning. The fear extinction network is well-characterized and includes the amygdala, hippocampus, and medial prefrontal cortex (mPFC; Quirk, Russo, Barron & Lebron, 2000; Corcoran & Maren, 2001; Santini, Ge, Ren, de Oritz, & Quirk, 2004; Bouton et al., 2006; Likhtik, Popa, Apergis-Schoute; Fidacaro, & Pare, 2008; Myers et al., 2011). Specifically, IL-mPFC is critical for the retrieval and consolidation of extinction of fear memories (Quirk et al., 2000; Sierra-Mercado, Padilla-Coreano, & Quirk, 2011) and extinction of drug seeking (Peters et al., 2008, Otis, Fitzgerald, & Mueller, 2014; Hafenbreidel, Rafa Todd, Twining, Tuscher, & Mueller, 2014). Because many of the same brain regions are recruited for the expression of fear- and drug-related extinction memories (Peters et al., 2009), literature on either type of memory could provide insight into the role of E₂ in extinction of cocaine seeking.

The role of E₂ in extinction of fear conditioning has been examined in humans as well as rodents. E₂ can enhance recall extinction of fear memories in both women and female rodents (Graham & Milad, 2013). This study was conducted in women who used hormonal contraceptives (HC) that contain estrogens and progesterone that inhibit ovulation by decreasing production of E₂ (Rivera, Yacobson, & Grimes, 1999). Importantly, combined HCs induce stable reductions in circulating E₂ compared to naturally cycling women during the follicular (high E₂) phase of the menstrual cycle (Pluchino et al., 2009). Women using HCs exhibited significantly poorer extinction recall compared to naturally cycling women as shown by higher skin conductance responses during fear conditioning and extinction. Moreover, administration of E₂ to naturally cycling women significantly reduced skin conductance responses and enhanced their ability to recall extinction memories (Graham & Milad, 2013). Extinction impairment was also observed in female rats treated with HCs, in which freezing behavior was used as an index to measure fear responses during extinction. HC-treated rats exhibited impaired extinction and continued to show elevated freezing responses in the presence of the CS compared to vehicle-treated rats. However, extinction impairment was reversed in HC-treated rats when serum E₂ levels were restored either by exogenous treatment of E₂ or by terminating HC treatment after fear conditioning (Graham & Milad, 2013). Additionally, systemic injections of ER β agonist (DPN) but not ER α agonist (PPT) prior to extinction training reduced freezing responses and facilitated extinction recall of fear in female rats (Zeidan et al., 2011). Thus, E₂ is necessary for extinction of fear memories in women and female rodents and this effect may be differentially regulated by ER β .

E₂ is also an important regulator of synaptic plasticity in females and males, in both rats and humans. Females have similar extinction recall to males when they undergo extinction during high E₂ phase of their menstrual cycle but have impaired extinction recall when they undergo

extinction during the low E₂ phase (Milad, Igoe, Lebron-Milad, & Novales, 2009, Milad et al., 2010). However, E₂ also plays an important role in the male brain. E₂ can be synthesized from circulating testosterone by the enzyme aromatase (Sharpe, 1998; Jones, Boon, Proietto, & Simpson, 2006; Gillies & McArthur, 2010). Inhibiting conversion to estrogens using an aromatase inhibitor significantly impaired extinction recall in male rats (Graham & Milad, 2014). Therefore, the effects of E₂ on extinction memories are not limited to females but include males as well.

The effects of E₂ on extinction of drug seeking has been examined in our labs. Similar to results observed following extinction of fear conditioning, systemic administration of E₂ facilitated extinction in a cocaine CPP paradigm (Twining, Tuscher, Doncheck, Frick, & Mueller, 2013). In this study, OVX rats were conditioned to associate a context with the rewarding effects of cocaine. During conditioning, rats received 2 days of E₂ treatment followed by 2 days of vehicle treatment to model naturally cycling rats. Systemic E₂ injections were either paired or explicitly unpaired with cocaine conditioning sessions. Another group of rats received E₂ daily during conditioning to test whether non-cycling groups exhibited similar acquisition or extinction patterns. Regardless of E₂ treatment during conditioning, all rats expressed a cocaine CPP which suggests that E₂ levels during conditioning did not affect subsequent CPP expression. Furthermore, during extinction, rats either received E₂ or vehicle prior to each extinction trial. Administration of E₂ to OVX rats promoted faster extinction compared to vehicle-treated rats. E₂-treated rats extinguished within a week, whereas vehicle-treated rats continued to perseverate for over 5 weeks. Importantly, extinction impairment in vehicle-treated rats was reversed by administration of E₂. Overall, this study has clear implications that E₂ may have therapeutic potential for women addicted to cocaine.

In summary, OVX female rats demonstrate impaired memory on a variety of behavioral tasks, as well as extinction of cocaine seeking, and these deficits can be rescued by E₂ replacement (Frick et al., 2010; Twining et al., 2013; Graham & Milad, 2013; Tuscher et al., 2015). E₂ can exert its actions via ER α and ER β which are expressed in the mPFC, amygdala, and hippocampus (Shughrue et al., 1998; Osterlund et al., 2000), structures involved in extinction learning and consolidation (Quirk & Mueller, 2008). Binding of E₂ to its receptors can stimulate intracellular signaling cascades, protein synthesis, and growth factor induction (Spencer et al., 2008; Frick, 2012; Frick et al., 2015) and potentially promote extinction through these mechanisms.

Effects of E₂ intrinsic excitability

The physiological mechanisms underlying the memory-enhancing effects of E₂ have been studied via electrophysiological recordings in hippocampal pyramidal cells. E₂ has the ability to enhance intrinsic excitability, which is defined by intrinsic membrane properties that affect how efficiently synaptic inputs are transformed into action potentials (APs; Wong & Moss, 1991; Kumar & Foster, 2002; Carrer, Araque, & Buno, 2003; Wu, Adelman, & Maylie, 2011). The hippocampus is necessary for associative learning and the degree of intrinsic excitability is positively correlated to the ability to learn hippocampus-dependent tasks (Thibault & Landfield, 1996; Moyer, Power, Thompson, & Disterhoft, 2000; Tombaugh, Rowe, & Rose, 2005; Wu et al., 2011). In agreement, an increase in excitability is reported following successful learning (Moyer, Thompson, & Disterhoft, 1996; Thompson, Moyer & Disterhoft, 1996).

Conversely, a lack of E₂ reduces excitability in hippocampal neurons. OVX rats that did not receive any E₂ treatment over a long period of time (~5 months) had reduced intrinsic membrane excitability of hippocampal pyramidal neurons compared to OVX controls that were

deprived of E₂ for 10 days (Wu et al., 2011). Slices from these rats required a larger depolarizing current to initiate firing compared to control OVX rats. Interestingly, bath-application of E₂ increased intrinsic excitability in the control OVX rats but not in the long-term E₂-deficient OVX rats. E₂ enhances excitability by suppressing the afterhyperpolarization (AHP) that follows action potentials. The AHP has multiple components, and is comprised of a fast AHP (fAHP), medium AHP (mAHP), and slow AHP (sAHP). Furthermore, the AHP acts as a negative feedback regulating excitability and spike frequency adaptation (Storm, 1990; Sah & Davies, 2000). E₂ can regulate the slow Ca²⁺-activated K⁺ current (*sI_{AHP}*) that mediates the sAHP in CA1 pyramidal neurons in hippocampal slices from OVX female rats (Kumar & Foster, 2002; Carrer et al., 2003). The *sI_{AHP}* in OVX rats was significantly larger compared to control rats, and bath-application of E₂ reduced the *sI_{AHP}* (Carrer et al., 2003). Overall, E₂ promotes excitability by suppressing *sI_{AHP}* in hippocampal neurons, which contributes to synaptic plasticity as well as learning and cognitive abilities. However, the role of E₂ in IL-mPFC neurons remains unknown.

Mechanisms underlying the role of E₂ in learning and memory

The molecular targets of E₂ are diverse but mounting evidence from clinical and pre-clinical studies suggest that Brain-derived neurotrophic factor (BDNF) may be a key mediator in regulating the memory-enhancing effects of E₂ (Singh, Meyer, & Simpkins, 1995; Aguirre & Baudry, 2009; Hill, 2012; Wu, Hill, Gogos, & VanDenBuuse, 2013; Fortress, Kim, Poole, Gould, & Frick, 2014). BDNF is an important regulator of neuroplasticity in different brain regions. BDNF is the most abundant neurotrophin in the central nervous system and has two receptors, its high-affinity receptor tropomyosin-related kinase B (TrkB) and the low-affinity receptor p75^{NTR}, which is a non-specific receptor for all neurotrophins (Chao, 2003; Huang & Reichardt, 2003; Reichardt, 2006). BDNF and its receptor TrkB play an important role in learning and memory.

Genetic and pharmacological disruptions of BDNF or its TrkB receptors have led to impairments in learning tasks. For example, heterozygous BDNF mutant mice were significantly impaired on spatial learning in a water maze test (Linnarsson, Bjorklund, & Ernfors, 1997). Spatial learning via the radial arm maze is positively correlated with enhanced BDNF mRNA as well as phosphorylated TrkB in the hippocampus (Mizuno, Yamada, Olariu, Nawa & Nabeshima, 2000; Mizuno, Yamada, He, Nakajima, & Nabeshima, 2003). Many of the trophic effects of E₂ and BDNF are similar, or even identical, which have led to the hypothesis that these signaling molecules can act synergistically to influence behavior, synaptic structure, and physiology of neurons.

Interactions between E₂ and BDNF

E₂ and BDNF are signaling molecules that can facilitate extinction of fear and drug seeking. As previously mentioned, exogenous treatment of E₂ enhanced extinction of fear conditioning in both female rats and humans (Zeidan et al., 2011; Graham & Milad, 2013). Facilitated extinction was also reported in a cocaine CPP paradigm when OVX female rats received systemic injections of E₂ (Twining et al., 2013). BDNF has been shown to play a similar role in extinction memory in male rats. For example, a single infusion of BDNF in IL-mPFC reduced conditioned fear (Peters, Dieppa-Perea, Melendez, & Quirk, 2010). Interestingly, infusions of BDNF into prelimbic mPFC, a structure involved in the expression of drug seeking behavior, were sufficient to suppress cocaine seeking behavior in a self-administration model. This type of suppression would normally occur only after extinction (Berglind et al., 2007). Additionally, a study by Otis et al., (2014) directly observed facilitated extinction of a cocaine CPP when TrkB receptors were potentiated by an infusion of BDNF in IL-mPFC of male rats. Extinction was impaired when BDNF activation was blocked by a highly selective TrkB receptor

antagonist, ANA-12 (Otis et al., 2014). Taken together, BDNF signaling in IL-mPFC regulates extinction of drug seeking and conditioned fear (Peters et al., 2010; Otis et al., 2014). However, the interaction of E₂ and BDNF in extinction of cocaine seeking in female rats is unclear.

An underlying mechanism of learning and memory involves dendritic plasticity. In general, E₂ is associated with greater dendritic spine density on pyramidal cells in the hippocampus and mPFC (Woolley & McEwen, 1993; Murphy, Cole, Greenberger, & Segal, 1998; McEwen & Elves, 1999; Christensen, Dewing, & Micevych, 2011; Tuscher, Luine, Frankfurt, & Frick, 2016). Similarly, BDNF/TrkB signaling has been shown to influence dendritic spine and synaptic density in hippocampal and cortical neurons (Gottmann, Mittmann, & Lessmann, 2009; Yoshii & Constantine-Paton, 2010; Luine & Frankfurt, 2013). Previous studies have reported that BDNF may be necessary for E₂ regulation of dendritic spines and ultimately synaptic transmission. E₂-mediated increases in dendritic spine density are attenuated by inhibiting TrkB receptors with K-252a in hippocampal slice cultures (Sato et al., 2007). Moreover, in aromatase knockout mice, there was a large decrease in both BDNF protein as well as dendritic spine density and this was rescued by treatment with estradiol benzoate (Sasahara et al., 2007). Thus, E₂ may regulate BDNF protein levels and spine density in the mPFC and hippocampus, regions crucial for extinction.

Regulation of dendritic morphology leads to the modulation of synaptic plasticity (Bourne & Harris, 2008). Previously, both BDNF and E₂ have been shown to be important modulators of synaptic plasticity by enhancing N-methyl-D-aspartate (NMDA) receptor mediated currents (Woolley, 1998; Foy et al., 1999; Poo, 2001; Lu, 2003; Cohen-Corey, Kidane, Shirkey, & Marshak, 2010; Otis et al., 2014). Furthermore, the effects of E₂ on synaptic transmission have been shown to be mediated by upregulation of BDNF (Scharfman, Mercurio, Goodman, Wilson,

& MacLusky, 2003). BDNF can have neurophysiological effects at different points of the estrous cycle, as BDNF expression is high during proestrus when E₂ levels are elevated compared to other phases of the cycle (Scharfman et al., 2003). In hippocampal slices, repetitive hilar stimuli evoked population spikes, inducing hyperexcitability in the CA3 region during the proestrus phase of the cycle. Stimulation to the CA3 region in slices derived from OVX rats did not evoke any spike responses. Furthermore, hyperexcitability in CA3 during proestrus was reversed by a non-specific Trk receptor antagonist, K-252a (Scharfman et al., 2003). Overall, this investigation suggests that E₂ may enhance synaptic transmission in hippocampal neurons via a BDNF-dependent mechanism.

Mechanisms implicated in synaptic plasticity and learning involve new protein synthesis and activation of cell-signaling cascades (Frick, 2012). Both E₂ and BDNF share common signal transduction pathways and transcription factors crucial for synaptic plasticity. These include signaling through the extracellular regulated protein kinase (ERK; Yamada & Nabeshima, 2003; Toran-Allerand et al., 1999; Boulware et al., 2013), the phosphatidylinositol 3-kinase (PI3-K; Znamensky et al., 2003; Mizuno et al., 2003; Fortress et al., 2013), Ca²⁺/Calmodulin-dependent protein kinase II (CaMKII; Sawai et al., 2002; Blanquet, Mariani, & Derer, 2003) and cAMP-response-element-binding protein (CREB; Ernfors & Bramham, 2003; McEwen, 2001; Boulware et al., 2005). Furthermore, E₂ regulates BDNF protein expression in a variety of brain regions. E₂ replacement in young adult OVX female rats enhances BDNF protein levels in the olfactory bulbs (Jeziarski & Sohrabji, 2000, 2001), hippocampus (Allen & McCarson, 2005; Gibbs 1998; Fortress et al., 2014), cortex (Allen & McCarson, 2005; Sohrabji, Miranda, & Toren-Allerand, 1995), amygdala (Liu et al., 2001; Zhou, Zhang, Cohen, & Pandey, 2005), septum (Gibbs 1999, Liu et al., 2001), dorsolateral area of the bed nucleus terminalis and the lateral habenular nucleus (Gibbs,

1999). E₂ has also shown to increase BDNF expression in the entorhinal cortex of OVX, aged female rats (Bimonte-Nelson, Nelson, & Granholm, 2004) as well as in the hippocampus of gonadectomized male mice (Solum & Handa, 2002). A possible explanation for the interaction between E₂ and BDNF may be due to the gene encoding BDNF, which contains a sequence similar to the ERE. This sequence is also found in other estrogen-regulated genes (Sohrabji et al., 1995). This suggests that the ER-ligand complex binds to the ERE-like motif on the BDNF gene and protects DNA from DNAase degradation, thereby regulating BDNF expression (Sohrabji et al., 1995).

A substantial amount of literature provides evidence that estrogenic regulation of BDNF is necessary for synaptic plasticity and memory (Srivastava et al., 2011; Luine & Frankfurt, 2013; Fortress et al., 2014). Regulation of BDNF by E₂ has been observed in brain structures such as the hippocampus and PFC, which are critical for memory formation (Luine & Frankfurt, 2013). Furthermore, low levels of BDNF and E₂ can be associated with anxiety-like behaviors (Gouveia et al., 2004), depression, (Russo-Neustadt & Chen, 2005; Marcus et al., 2005) and impaired memory formation (Matsushita et al., 2005; Bian, Zhang, Zhang, & Zhao, 2005). Specifically, both E₂ and BDNF have facilitated extinction of a cocaine CPP, independently (Twining et al., 2013; Otis et al., 2014). Therefore, this suggests that the regulation of BDNF by E₂ may play an important role in extinction of cocaine seeking.

Clinical relevance

Drug-associated memories can be strengthened by environmental stimuli (cues) that promote compulsive drug taking and trigger relapse (Carter & Tiffany, 1999; Shalev, Grimm, & Shaham, 2002; See, 2005). Both clinical and preclinical studies of substance use disorders suggest that females are more vulnerable than males to addiction, including initiation of drug taking,

bingeing, and relapse (Anker & Carroll, 2010). Treatment of addictive disorders can be modeled through extinction and has been shown to be effective in preclinical investigations. Specifically, E₂ facilitates extinction in female rodents (Zeidan et al., 2011; Twining et al., 2013; Graham & Milad, 2013), therefore, it may be more clinically relevant to investigate mechanisms underlying E₂-induced facilitation of extinction. Furthermore, pharmacological activation of BDNF or E₂ signaling would enhance extinction-based therapies for learning and memory related disorders, such as addiction.

AIMS

Aim 1

In the present study, we aimed to investigate the mechanisms mediating E₂-induced facilitation of extinction of cocaine seeking. Data from our lab have shown that E₂ is necessary for extinction of a cocaine-induced CPP and lack of it results in persistent deficits of extinction (Twining et al., 2013). The goal of this study was to target BDNF/TrkB signaling as a potential mechanism for E₂-induced enhancement of extinction as direct infusions of BDNF in the IL-mPFC has shown to facilitate extinction in male rats (Otis et al., 2014). To date, this mechanism has not been examined in female rats. Our hypothesis is that E₂ will facilitate extinction of cocaine seeking and this effect will be impaired by systemic or IL-mPFC infusions of Trk receptor antagonists. Since we aimed to impair acquisition of extinction, pharmacological manipulations were made prior to extinction trials.

Aim 2

In aim 2, we determined the interactions between BDNF and E₂ via electrophysiological recordings in IL-mPFC neurons derived from female OVX rats. We measured intrinsic

excitability which lowers the threshold for synaptic changes and is a neural correlate of learning. A significant amount of attention has been given to synaptic plasticity as a mechanism for learning. However, recent studies have demonstrated that memory storage cannot be exclusively based on a synaptic model and that intrinsic excitability plays a critical role in adaptive behaviors (Frick & Johnston, 2005; Song, Detert, Sehgal, & Moyer, 2012; Zhang & Linden, 2003), such as extinction. E₂ enhances intrinsic physiology in hippocampal neurons (Kumar & Foster, 2002; Carrer et al., 2003) but the effects of E₂ on IL-mPFC neurons remain unknown. Furthermore, there is no known data that BDNF influences intrinsic excitability in the hippocampus or any other regions necessary for extinction. This study aimed to examine whether E₂ enhances intrinsic excitability and if this effect was dependent on BDNF/TrkB signaling. To test this, OVX female rats received systemic injections of E₂ or vehicle to determine if priming with E₂ affected membrane properties as previously shown (Schiess, Joels, & Shinnick-Gallagher, 1988). Rats were euthanized and patch-clamp recordings were conducted in layer 5 of IL-mPFC pyramidal neurons. Throughout recordings, brain slices were incubated with vehicle, E₂ (25 nM), vehicle + K-252a, E₂ + K-252a and excitability was measured. We hypothesized that E₂ would enhance intrinsic excitability of IL-mPFC pyramidal neurons, and that blockade of Trk receptors would attenuate the effects of E₂.

METHODS

Subjects and Surgery

Female Long-Evans rats weighing between 275-300 g were individually housed in clear plastic cages. Rats were maintained on a 14-h light/10-h dark cycle and had unlimited access to water and standard laboratory chow (Teklad, Harlan Laboratories). Rats were weighed and handled daily for approximately 3 days prior to surgery and before the start of experiments. All experimental protocols were approved by the Institutional Animal Care and Use Committee at the University of Wisconsin-Milwaukee in accordance with National Institute of Health guidelines.

Surgeries were performed as previously described (Frick et al., 2004; Twining et al., 2013; Otis, Kidane, & Mueller, 2013). Rats were anesthetized with a mixture of ketamine/xylazine (90mg/kg/10.5mg/kg, i.p.) and underwent bilateral OVX using a dorsal approach (Frick et al., 2004). A single, horizontal incision was made along the spine and the ovary was isolated. The tip of the uterus was clamped and ligated and the ovary was removed with a scalpel. The remaining tissue was returned to the abdomen. The same procedure was repeated on the other ovary, and the incision was closed with sterile sutures and wound clips. Next, rats were implanted with a double-barrel guide cannula aimed bilaterally at IL-mPFC (anteriorposterior, +2.8; mediolateral \pm 0.6, and dorsoventral, -4.4 mm relative to bregma). Following surgeries, rats were given an antibiotic (penicillin G procaine, 75,000 units in 0.25 mL) and an analgesic (carprofen, 5.0 mg in 0.1 mL) subcutaneously and then allowed to recover for approximately 7 days before behavioral testing. After completion of behavioral procedures, verification of cannula placements was conducted using a cresyl violet stain.

Drugs

Cocaine HCl (National Institute on Drug Abuse) was dissolved in sterile 0.9% saline at a concentration of 10 mg/mL, and administered systemically at a dose of 10 mg/kg, i.p. To ensure that E₂ levels did not build over time from repeated infusions, a water-soluble form of E₂ dissolved in 2-hydroxypropyl- β -cyclodextrin (HBC) that is metabolized within 24h was used (Pitha & Pitha, 1985). HBC vehicle and HBC-encapsulated E₂ (Sigma Aldrich) were dissolved in sterile 0.9% saline (0.2 mg/mL) and injected i.p. at a dose of 0.2 mg/kg (Gresack & Frick, 2006). ANA-12 (Sigma Aldrich) was dissolved in 1% DMSO in physiological saline (Zhang et al., 2015) and administered i.p. at 0.5 mg/kg, i.p. (Cazorla et al., 2011). K-252a (Sigma Aldrich) was dissolved in vehicle of 50% DMSO in artificial cerebral spinal fluid (aCSF) and infused into the IL-mPFC at 3.5 ng/0.3 μ l/side (50 μ M; Xin et al., 2014). For electrophysiological recordings, 25 nM of β -estradiol (not HBC encapsulated) was dissolved in 100% DMSO and diluted with aCSF to a final DMSO concentration of 0.0001%. K-252a was dissolved in 100% DMSO and bath-applied at 100 nM and diluted to a final concentration of 0.001% DMSO (Montalbano, Baj, Papadia, Tongiorgi, & Sciancalepore, 2013).

Conditioned place preference

Place preference apparatus

Testing and conditioning were conducted in a 3-chamber apparatus in which 2 larger conditioning chambers (13" x 9" x 11.5") were separated by a smaller chamber (6" x 7" x 11.5"). The larger conditioning chambers had wire mesh flooring with white walls, whereas the other had gold-grated flooring with a black wall. The center chamber had aluminum sheeting as flooring. All floors were raised 1.5", with removable trays placed beneath. Removable partitions were

used to isolate the rats within specific chambers during conditioning. During baseline and CPP trials, the doors were removed to allow free access to the entire apparatus. Each of the larger chambers contain two infrared photobeams separated by 3". If the beam furthest from the door was broken, then the rat was determined to be in the larger chamber. If only the beam closest to the center chamber was broken, then the rat was determined to be in the center chamber. During all phases of the experiments the room was kept in semi-darkness.

Conditioning and Testing

A pre-test determined baseline preferences by placing the rats into the center chamber with free access to the entire apparatus for 15 min and recording time in each chamber. Rats spent an equal amount of time in the larger conditioning chambers, but less time in the center chamber. ANOVA revealed an effect of chamber for all rats during baseline test ($F_{(2,204)} = 98.73, p < 0.001$), and *post-hoc* analyses confirmed that less time was spent in the center chamber than either of the conditioning chambers ($p < 0.001$). Therefore, an unbiased procedure was used, in which rats were randomly assigned to receive cocaine in one of the two larger chambers, independent of baseline preference scores. After a pre-test, rats were conditioned to associate one chamber, but not another, with cocaine in a counterbalanced fashion over 8 days. Injections were given immediately before placing the rats in their chambers for 20 min conditioning sessions.

Following conditioning, rats went through extinction training in which they were placed into the center chamber and allowed free access to the entire apparatus. In general, when the hypothesis was drug-induced facilitation of extinction, CPP extinction trials lasted for 15 min. Alternatively, when the hypothesis was drug-induced impairment of extinction, CPP extinction trials lasted for 30 min in length.

Experiment one

All rats received daily 0.2mg/kg (Gresack & Frick, 2006) i.p. injections of E₂ throughout conditioning phase. Rats received E₂ treatment one hour before conditioning trials which lasted for 8 days. Conditioning trials consisted of 4 pairings with cocaine and 4 pairings with saline. Following conditioning, rats did not receive any hormone treatment for 2 days before CPP testing (Twining et al., 2013). To determine whether K-252a impairs extinction learning of a cocaine CPP, rats received E₂ injections one hour prior to a CPP trial and were infused with 50 µM of K-252a (Xin et al., 2014) or aCSF vehicle 15 min prior to a CPP trial (30 min). K-252a was infused 15 min prior to a CPP trial as previous studies have shown that infusions of K-252a in the IL-mPFC approximately 15 min prior to testing impaired behavioral performance in other paradigms (Xin et al., 2014). To control for hormonal manipulations, rats received HBC one hour prior to a CPP trial and were also infused with K-252a or aCSF 15 min prior to a CPP trial (30 min). A 30 min time period was chosen for the extinction trials to observe impairment as extinction proceeds faster in controls.

Experiment two

We then examined whether direct inactivation of TrkB receptors impairs extinction by systemically administering the selective TrkB receptor antagonist, ANA-12. One hour prior to each CPP trial (15 min), rats received either systemic injections of 0.2 mg/kg E₂ (Gresack & Frick, 2006) and 0.5 mg/kg ANA-12 (Cazorla et al., 2011; Zhang et al., 2015) or E₂ and vehicle. The dose for ANA-12 was chosen as this dose impaired behavioral performance in other paradigms (Zhang et al., 2015). ANA-12 was systemically administered 1 hour prior to a CPP trial because active concentrations have been detected as early as 30 minutes and up to 6 hours after systemic injections (Cazorla et al., 2011). A 15 min time period was chosen for extinction

trials in this experiment even though the hypothesis was to impair extinction because 30 min extinction trials produced large variability in previous experiments.

Experiment three

As previously mentioned, experimental manipulations of extinction can be made prior to extinction trials, whereas consolidation of extinction can be manipulated immediately following extinction training (Millan et al., 2011). Therefore, to determine whether systemic administration of E₂ is necessary for consolidation of extinction, rats received systemic injections of 0.2 mg/kg E₂ (Gresack & Frick, 2006) or HBC vehicle immediately after daily CPP extinction trials (15 min). Rats were given shortened 15 min extinction trials to limit extinction across trials, as we hypothesized that E₂ would enhance the consolidation of extinction of a cocaine-induced CPP.

Histology

After behavioral testing, rats were euthanized with an overdose of ketamine and perfused with 0.9% saline followed by 10% buffered formalin phosphate. Brains were removed and placed in 30% sucrose/formalin solution. Following brain submersion, 40 μM thick coronal sections were sliced using a microtome from brain regions in which cannula were implanted. Sections were then mounted and stained with cresyl violet. Injector tip locations were confirmed using a rat brain atlas (Paxinos & Watson, 2007).

Data analysis

Drug seeking behavior was analyzed using a three-way ANOVA to compare time within each chamber across trials and between groups (Twining et al., 2013; Otis et al., 2014). When appropriate main interaction effects were detected, Fisher's LSD post hoc tests were used to make pairwise comparisons.

Patch-clamp electrophysiology

Female rats aged 3 months were OVXed and allowed to recover for 4-5 days. Rats received systemic injections of E₂ or HBC for three consecutive days before being euthanized for patch-clamp recordings. They were anesthetized with isoflurane, and their brains were rapidly removed and transferred to ice-cold, oxygenated (95%/O₂/5%CO₂) aCSF containing the following composition (in mM): 124 NaCl, 2.8 KCl, 1.25 NaH₂PO₄, 2 MgSO₄, 2 CaCl₂, 26 NaHCO₃, and 20 dextrose. Coronal slices were cut 400 μ m in ice-cold aCSF using a vibrating blade microtome (Leica VT1200). Slices were allowed to recover in warm aCSF (32-35°C) for 30 min followed by incubation in room-temperature aCSF for the remainder of the experiment (until the end of the day). Slices were transferred to a submersion chamber, mounted, and perfused with aCSF (~2 ml/min; 32°C). Pyramidal neurons with visible apical dendrites in layer V of the IL-mPFC were visualized with differential interference contrast using a 60X water-immersion lens on an upright Eclipse FN1 microscope (Nikon Instruments). Whole-cell patch recordings of IL-mPFC pyramidal neurons were obtained using fire polished borosilicate glass pipettes (3-8 M Ω), filled with internal solution containing the following (in mM): 110 K-gluconate, 20 KCL, 10 HEPES, 2 MgCl₂, 2 ATP, 0.3 GTP, 10 phosphocreatine; 0.2% biocytin, pH 7.3, and 290 mOsm. Intrinsic excitability were obtained with current clamp using a MultiClamp 700B (Molecular Devices) patch-clamp amplifier connected to a Digidata 1440A digitizer (Molecular Devices). The liquid-liquid junction potential (measured as 13 mV) was compensated for all recordings. All electrophysiological data were analyzed using Clampfit (Molecular Devices).

Experimental Design

After 10 min of stable recordings, neurons were polarized to approximately -60 mV to control for variance in resting membrane potential. A series of 1s current steps were applied (-40

to 500 pA; 10 pA steps) and the number of evoked APs were recorded to measure basic membrane properties such as input resistance and rheobase. To measure input resistance, current pulses were injected and the resulting voltage deflections were measured to create an IV plot. Furthermore, a rheobase was measured which is the minimum amount of current required to elicit a single action potential. Rheobase was analyzed in a subset of pyramidal neurons by applying 1s current steps with 10 pA increments until a single AP is elicited. Intrinsic excitability was measured by applying a 2s depolarizing step every 7.5s and evoked APs were recorded. The level of depolarizing step was adjusted to rheobase and remained constant throughout the experiment (Otis et al., 2013). To measure effects of E₂ on intrinsic excitability, 25 nM of E₂ was bath-applied and baseline current steps were applied every 5 min for approximately 30 min. To ensure that excitability was enhanced by E₂ and not by the current pulses on its own, the same protocols were repeated for the same amount of time in slices that remained in aCSF. To measure effects of K-252a on E₂ induced excitability, slices were bathed in 100 nM K-252a (Montalbano et al., 2013) for approximately 20 min before bath-application of E₂. Following recording, brain slices were fixed in phosphate-buffered formalin overnight.

Immunohistochemistry

To confirm that patch-clamp recordings were from IL-mPFC neurons, biocytin-filled pyramidal neurons were washed in 0.1 M PBS, followed by 1% NaBH₄ in PBS and 10% normal goat serum (NGS). The slices incubated overnight with 3% NGS, 0.2% Triton-X, and PBS. After 24 hours, slices were washed in PBS and incubated for 2h with a green fluorescent antibody (streptavidin, 1:250; Invitrogen). Slices were washed with PBS before being mounted with antifade mounting medium and coverslipped. Slices were visualized using 20X magnification with green fluorescent light, to locate neurons and verify that they were pyramidal.

Electrophysiological data analysis

Electrophysiological data were analyzed using Clampfit (Molecular Devices). Basic neuron properties were examined: input resistance, resting membrane potential, AP half-width, and AP amplitude. To analyze slow afterhyperpolarization (sAHP), voltage was recorded 1s following current offset which was then subtracted from baseline voltage before current injection (Kaczorowski, Davis, & Moyer, 2012). Basic measures of intrinsic excitability were analyzed before and immediately following drug application. Finally, to analyze AP frequency, the average number of spikes was plotted against time.

RESULTS

Trk receptor blockade in IL-mPFC facilitated extinction of a cocaine CPP

Previous work conducted in our lab has shown that E₂, when administered daily, facilitates extinction of cocaine seeking in female rats (Twining et al., 2013). The mechanism underlying E₂-induced facilitation of extinction, however, is unknown. BDNF has a role in regulating the memory-enhancing effects of E₂ in other paradigms (Singh et al., 1995; Aguirre & Baudry, 2009; Fortress et al., 2014) and facilitates extinction of a cocaine CPP in male rats (Otis et al., 2014). BDNF targets a Trk receptor, and therefore we determined the necessity of Trk receptor activation for E₂-induced facilitation of extinction in OVX female rats. We targeted Trk receptors using a non-selective antagonist, K-252a. We hypothesized that K-252a would prevent the E₂-induced facilitation of extinction. For this experiment, rats were surgically implanted with bilateral guide cannula aimed at IL-mPFC (for all IL-mPFC sites, see Figure 1A). Rats were pretested, conditioned, and subjected to daily CPP trials. Rats were given lengthened extinction trials (30 min) to promote learning, as we hypothesized that K-252a would impair extinction (Otis et al., 2014). One hour prior to each CPP trial, all OVX female rats received 0.2 mg/kg of E₂ (Gresack & Frick, 2006) systemically to facilitate extinction of a cocaine CPP (Twining et al., 2013). Rats were given a single bilateral IL-mPFC infusion of K-252a (n = 8; 3.5 ng/0.3 μ l/side; Xin et al., 2014) or aCSF vehicle (n = 8) 15 min prior to a 30 min extinction trial (Figure 1B). A mixed model three-way ANOVA revealed a significant chamber by trial by group interaction ($F_{(10, 140)} = 2.502, p < 0.05$), and subsequent two-way ANOVA indicated a chamber by trial interaction for K-252a-treated rats ($F_{(10, 70)} = 5.961, p < 0.001$), but not vehicle-treated rats ($F_{(10, 70)} = 0.882, p > 0.05$).

Post hoc analysis confirmed that both groups spent significantly more time in the previously cocaine-paired chamber during the first CPP trial ($p < 0.05$), indicating that both groups expressed a CPP. During subsequent extinction trials (i.e., trials 2,3,5, and 6), aCSF-infused rats continued to spend significantly more time in the cocaine-paired chamber than in the saline-paired chamber (*post hoc* $p < 0.05$), while K-252a-treated rats spent an equivalent amount of time in the cocaine-paired and saline-paired chambers during extinction trials 2,3,4, and 6 (*post hoc* $p > 0.05$). Overall, we did not observe the expected E₂-induced facilitation of extinction shown previously by our lab (Twining et al., 2013), and contrary to expectations K-252a facilitated extinction of a cocaine CPP.

We repeated the experiment, and extended the infusion protocol to include daily infusions of K-252a across four days instead of one. Previous work in our lab has demonstrated that four daily systemic injections or infusions into IL-mPFC of NMDA receptor antagonists impaired extinction (Otis et al., 2014; Hafenbreidel et al., 2014). In addition to systemic injections of 0.2 mg/kg E₂ (Gresack & Frick, 2006), rats received 3.5 ng/0.3 μ l/side of K-252a ($n = 8$; Xin et al., 2014) infusions or vehicle aCSF ($n = 10$) into IL-mPFC (Figure 2A for all IL-mPFC infusion sites) 15 min prior to each of the first four extinction trials (30 min; Figure 2B). Rats did not receive any infusions during subsequent extinction trials (i.e., 5 and 6; Otis et al., 2014). ANOVA revealed no significant chamber by trial by group interaction ($F_{(10, 160)} = 0.761, p > 0.05$). However, there was a significant chamber by trial interaction, ($F_{(10, 160)} = 2.768, p < 0.05$), and a significant effect of chamber overall ($F_{(2, 32)} = 35.613, p < 0.001$). *Post hoc* analysis confirmed that both K-252a-treated and aCSF-treated rats spent more time in the previously cocaine-paired chamber than in the saline-paired chamber during the first trial ($p < 0.05$). However, K-252a-treated rats spent equivalent time in the cocaine-paired and saline-paired chambers (*post hoc* $p >$

0.05) during all subsequent trials (i.e. 2-6). aCSF-treated rats spent significantly more time in the cocaine-paired chambers during trials 2 and 4. Although E₂-treated rats did extinguish across days, inhibition of Trk receptors did not impair this effect and rather appeared to facilitate extinction. Thus, at the dose used in this study, Trk receptors in the IL-mPFC do not mediate E₂-induced facilitation of extinction of cocaine seeking.

Figure 1




Figure 1. A single IL-mPFC infusion of non-selective Trk receptor antagonist facilitated extinction of a cocaine CPP. (A) Coronal drawings (bregma, +3.72 mm; Paxinos & Watson, 2007; Copyright 2007) showing placements of injector tips for K-252a (n = 8) and vehicle (n = 8). Following conditioning IL-mPFC infusion (arrow) of (B) Trk receptor antagonist, K-252a prior to the first CPP trial facilitated extinction. *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$.

Figure 2





Figure 2. Multiple IL-mPFC infusions of non-selective Trk receptor antagonist facilitated extinction of a cocaine CPP. (A) Coronal drawings (bregma, +3.72 mm; Paxinos & Watson, 2007; Copyright 2007) showing placements of injector tip placements for K-252a (n = 8) and vehicle (n = 10). Following conditioning IL-mPFC infusions (arrows) of (B) Trk receptor antagonist, K-252a prior to the first four CPP trials facilitated extinction. *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$.

Trk receptor blockade in IL-mPFC did not impair extinction of a cocaine CPP in the absence of E₂

Trk receptor blockade did not impair extinction when rats were administered E₂ prior to each CPP trial but facilitated extinction of a cocaine CPP. To control for hormonal manipulations, we investigated whether Trk receptor antagonist infusions into IL-mPFC would block extinction in the absence of E₂. Following conditioning, all OVX female rats received HBC vehicle one hour prior to a CPP trial (Twining et al., 2013). Rats were given a single bilateral IL-mPFC infusion of K-252a (n = 6; 3.5 ng/0.3 µl/side; Xin et al., 2014) or aCSF vehicle (n = 6) 15 min prior to a 30 min extinction trial (Figure 3). A mixed model three-way ANOVA revealed a significant effect of chamber overall ($F_{(2, 20)} = 26.06, p < 0.001$) and a significant chamber by trial interaction ($F_{(2, 80)} = 3.206, p < 0.01$). However, there was no significant group by chamber by trial interaction ($F_{(8, 80)} = 1.840, p > 0.05$). *Post hoc* analysis confirmed that both vehicle-treated and K-252a-treated rats showed a significant preference for the cocaine paired chamber on trial 1 (*post hoc* $p < 0.05$). However, both groups did not show a significant preference on all subsequent trials (*post hoc* $p > 0.05$). Inconsistent with previous findings (Twining et al., 2013), HBC-treated rats did show evidence of extinction, although extinction training was not extended for the 16-32 day procedure used previously in which CPP expression was initially poor but stabilized across days. In addition, blocking Trk receptors in IL-mPFC at the selected dose did not impair extinction of a cocaine CPP.

Figure 3

A



B




Figure 3. A single IL-mPFC infusion of non-selective Trk receptor antagonist did not disrupt extinction of a cocaine CPP in the absence of E₂ (A) Coronal drawings (bregma, +3.72 mm; Paxinos & Watson, 2007; Copyright 2007) showing placements of injector tips for K-252a (n = 6) and vehicle (n = 6). Following conditioning IL-mPFC infusion (arrow) of (B) Trk receptor antagonist, K-252a prior to the first CPP trial had no effect on extinction.

** $p < 0.01$

E₂-induced facilitation of a cocaine CPP was disrupted by systemic injections of TrkB receptor antagonist

The previous experiments indicate that an infusion of the non-selective receptor antagonist, K-252a, in IL-mPFC did not impair extinction of a cocaine CPP. This suggested that Trk receptors in the IL-mPFC may not play a role in mediating the effects of E₂ in the enhancement of extinction. Thus, we examined whether TrkB receptors mediated extinction by using systemic injections of an antagonist. K-252a does not cross the blood-brain barrier, so we used a selective TrkB receptor antagonist, ANA-12, that is known to cross the blood-brain barrier effectively (Cazorla et al., 2011; Zhang et al., 2015). Second, in the experiments that we conducted so far (Figure 1, Figure 2, Figure 3), we reported a large amount of variability in the vehicle-treated rats with 30 min extinction trials and this variability was not observed previously in male rats (Otis et al., 2014). Furthermore, E₂-induced enhancement of extinction in female rats was reported with 15 min extinction trials (Twining et al., 2013). Therefore, for this experiment, 15 min extinction trials were used, which is consistent with previous data collected from OVX female rats (Twining et al., 2013). All rats received 0.2 mg/kg of E₂ (Gresack & Frick, 2006) one hour prior to each CPP trial. In addition to E₂ injections, either rats received a systemic injection of 0.5 mg/kg of ANA-12 (n = 9; Cazorla et al., 2011; Zhang et al., 2015) or vehicle (n = 10) one hour before a CPP trial (Figure 4). This dose has been reported to impair performance in a number of behavioral paradigms (Zhang et al., 2015). Furthermore, rats received systemic injections of ANA-12 one hour prior to a CPP trial as active concentrations of ANA-12 have been detected as early as 30 minutes and up to 6 hours after an injection (Cazorla et al., 2011). ANOVA revealed no significant trial by chamber by group interaction ($F_{(10, 170)} = 0.1203, p > 0.05$). However, there was a significant effect of trial by chamber ($F_{(10, 170)} = 3.926, p < 0.001$) and

an overall effect of chamber ($F_{(2, 34)} = 38.457, p < 0.001$). *Post hoc* analysis confirmed that both ANA-12-treated and vehicle-treated rats spent more time in the previously cocaine-paired chamber than in the saline-paired chamber during the first trial ($p < 0.001$). However, vehicle-treated rats did not show a significant preference for the cocaine-paired chamber during subsequent trials (*post hoc* $p > 0.05$). ANA-12-treated rats showed a significant preference for the cocaine-paired chamber during trials 2, 3, and 5 (*post hoc* $p < 0.05$). Previous experiments in this particular study did not report an E_2 -induced facilitation of extinction consistently. However, in this experiment, E_2 -facilitated extinction appears to be impaired by TrkB receptor antagonist, ANA-12.

Figure 4




Figure 4. Injections of selective TrkB receptor antagonist disrupted E_2 -induced facilitation of extinction of a cocaine CPP. Systemic injections (arrows) of ANA-12 ($n = 9$) or vehicle ($n = 10$) before the first 4 CPP trials impaired E_2 -induced facilitation of extinction. *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$.

E₂ potentiates IL-mPFC pyramidal neuron excitability

Intrinsic excitability of hippocampal pyramidal neurons is potentiated by E₂ by suppressing sAHP following action potentials (Kumar & Foster, 2002; Carrer et al., 2003; Wu et al., 2011). E₂ can enhance intrinsic excitability, which is an underlying cellular mechanism of successful learning (Moyer et al., 1996; Thompson et al., 1996) and may offer a potential mechanism for extinction. To date, whether E₂ enhances intrinsic excitability in the IL-mPFC pyramidal neurons remains unknown. To test this, female OVX rats received systemic injections of E₂ for three days before they were euthanized for recordings to ensure the effects of excitability with E₂ priming. Previously, E₂ priming has been shown to affect membrane properties and excitability in neurons (Schiess et al., 1988). Furthermore, we used patch-clamp electrophysiology to record APs evoked by brief current pulses in layer 5 IL-mPFC pyramidal neurons (Figure 5A). Bath-application of E₂ (25 nM) increased the number of evoked APs in slices extracted from rats that had previously received systemic injections of E₂ (Figure 5C). To ensure that excitability is drug-dependent and not due to the current injections, cells in the control group (n = 8) received the same number of current injections over time as the experimental group (n = 9; Figure 5C). ANOVA revealed a significant effect of treatment ($F_{(2,105)} = 8.139, p < 0.001$), an effect of time ($F_{(5, 105)} = 2.511, p < 0.05$) but no treatment by time interaction ($F_{(5, 105)} = 1.527, p > 0.05$). Furthermore, E₂ caused membrane depolarization and decreased rheobase, which are also indicative of increased excitability and this was not observed in control slices treated with aCSF (Otis et al., 2013; Table 1). E₂ did not reduce sAHP which suggests that E₂ does not increase excitability by reducing sAHP in IL-mPFC pyramidal neurons.

Figure 5




Figure 5. Bath-application of E₂ potentiates IL-mPFC pyramidal neuron excitability in slices from OVX rats that were primed with E₂. (A) Photomicrograph of a biocytin-filled pyramidal neuron. (B) Individual traces of current-evoked APs from an IL-mPFC pyramidal neuron before and after E₂ application. (C) E₂ increases AP frequency. *** $p < 0.001$. Error bars indicate SEM.

Table 1. Effects of E₂ on intrinsic excitability of IL-mPFC pyramidal neurons in slices from OVX rats primed with E₂

Drug	Time	R _N (MΩ)	V _m (mV)	Rheo (pA)	sAHP (mV)	AP _{hw} (ms)	AP _{amp} (mV)
E ₂	Pre	277 ± 35	-60 ± 0.6	33 ± 6	0.15 ± 0.3	1.25 ± 0.1	38.8 ± 2.3
	Post	318 ± 44*	-59 ± 0.0*	27 ± 6*	-0.01 ± 0.2	1.20 ± 0.1*	33.8 ± 2.6
aCSF	Pre	347 ± 30	-59 ± 1	24 ± 3	-0.8 ± 0.3	1.50 ± 0.1	39 ± 1.3
	Post	403 ± 44**	-59 ± 0.8	23 ± 4	-0.08 ± 0.2	1.50 ± 0.1	33 ± 1.4*

E₂, 17β-estradiol; R_N, input resistance; V_m, resting potential; Rheo, rheobase; sAHP, slow afterhyperpolarization; AP_{hw}, Action Potential half-width; AP_{amp}, Action Potential amplitude. **p* < 0.05, ***p* < 0.001 as compared to pre-E₂ application.

To control for hormonal manipulations prior to recording, we systemically injected OVX female rats with HBC vehicle for three days before they were euthanized for patch-clamp recordings. The brain slices extracted from these rats were incubated in 25nM E₂ or aCSF and the number of evoked APs were recorded from IL-mPFC (Figure 6A). To ensure that excitability is drug-dependent and not due to the current injections, cells in the control group (n = 9) received the same number of current injections over time as the experimental group (n = 9; Figure 6C). Bath-application of E₂ (25 nM) increased the number of evoked APs in slices extracted from rats that previously received HBC injections (Figure 6C). ANOVA revealed an effect of treatment ($F_{(2,112)} = 14.976, p < 0.001$), but no effect of time ($F_{(5,112)} = 1.916, p > 0.05$) or treatment by time interaction ($F_{(5,112)} = 0.263, p > 0.05$). E₂ did not cause membrane depolarization or reduce sAHP in IL-mPFC neurons (Table 2). Thus, bath-application of E₂ enhances excitability in IL-mPFC pyramidal neurons without E₂ priming.

Figure 6




Figure 6. Bath-application of E₂ potentiates IL-mPFC pyramidal neuron excitability in slices from OVX rats that were not primed with E₂. (A) Photomicrograph of a biocytin-filled pyramidal neuron. (B) Individual traces of current-evoked APs from an IL-mPFC pyramidal neuron before and after E₂ application. (C) E₂ increases AP frequency. *** $p < 0.001$. Error bars indicate SEM.

Table 2. Effects of E₂ on intrinsic excitability of IL-mPFC pyramidal neurons in slices from OVX rats that were not primed with E₂

Drug	Time	R _N (MΩ)	V _m (mV)	Rheo (pA)	sAHP (mV)	AP _{hw} (ms)	AP _{amp} (mV)
E ₂	Pre	273 ± 38	-59 ± 0.5	29 ± 4	-0.03 ± 0.1	1.0 ± 0.0	38.9 ± 2.9
	Post	306 ± 44*	-58 ± 0.5	19 ± 4**	-0.1 ± 0.2	0.95 ± 0.0**	33.9 ± 3.0*
aCSF	Pre	309 ± 52	-57 ± 0.5	36 ± 7	-0.4 ± 0.2	1.1 ± 0.0	34.6 ± 2.7
	Post	310 ± 21	-57 ± 0.8	28 ± 6**	-0.1 ± 0.2	1.07 ± 0.0**	30.6 ± 3.1*

E₂, 17β-estradiol; R_N, input resistance; V_m, resting potential; Rheo, rheobase; sAHP, slow afterhyperpolarization; AP_{hw}, Action Potential half-width; AP_{amp}, Action Potential amplitude. * $p < 0.05$, ** $p < 0.001$ as compared to pre-E₂ application.

Trk receptor blockade prevented E₂-induced potentiation of IL-mPFC pyramidal neuron excitability

Intrinsic excitability of cortical pyramidal neurons is potentiated by Trk receptor activation (Marongiu, Imbrosci, & Mittmann, 2013) and may offer a potential a mechanism by which Trk receptors mediate extinction. Whether Trk receptor blockade prevents E₂-induced enhancement of intrinsic excitability in IL-mPFC pyramidal neurons is unknown. Bath-application of E₂ enhances excitability in IL-mPFC neurons with or without prior systemic E₂ administration (Figure 5, Figure 6). To examine the role of Trk receptors in E₂-induced excitability of IL-mPFC neurons (Figure 7A), OVX female rats received systemic HBC vehicle injections before they were euthanized for patch-clamp recordings. Slices were incubated for 20 min in 100 nM K-252a (Montalbano et al., 2013) and E₂ (25 nM) was bath-applied. The effect of E₂ was blocked by bath-application of K-252a (Figure 7B), indicating that Trk receptor blockade prevents E₂-induced potentiation of IL-mPFC neuron excitability. Comparing neurons treated with E₂ (n = 9), E₂ + K-252a (n = 7), and control (n = 9) ANOVA revealed an effect of treatment ($F_{(4,147)} = 4.932, p < 0.001$), but no effect of time ($F_{(5,147)} = 1.750, p > 0.05$), or treatment by time interaction ($F_{(10,112)} = 1.061, p > 0.05$). E₂ reduces rheobase, AP half-width, and AP amplitude and this change is not present in the presence of K-252a (Table 3). Thus, E₂ potentiates excitability in IL-mPFC neurons via a Trk receptor-dependent mechanism.

Figure 7




Figure 7. Blockade of Trk receptors prevent E₂-induced potentiation of IL-mPFC neuronal intrinsic excitability (A) Photomicrograph of a biocytin-filled pyramidal neuron. (B) Trk receptor blockade prevents E₂-induced increases in AP. * $p < 0.001$. Error bars indicate SEM.**

Table 3. Effects of E₂ and K-252a on intrinsic excitability of IL-mPFC pyramidal neurons

Drug	Time	R _N (MΩ)	V _m (mV)	Rheo (pA)	AP _{hw} (ms)	AP _{amp} (mV)
E ₂	Pre	273 ± 38	-59 ± 0.5	29 ± 4	1.0 ± 0.0	38.9 ± 2.9
	Post	306 ± 44*	-58 ± 0.5	19 ± 4**	0.95 ± 0.0**	33.9 ± 3.0*
E ₂ +K-252a	Pre	266 ± 39	-59 ± 0.1	40 ± 4	1.5 ± 0.1	32.9 ± 4.0
	Post	322 ± 38*	-60 ± 0.4	37 ± 6	1.4 ± 0.05	29.1 ± 4.1

E₂, 17β-estradiol; R_N, input resistance; V_m, resting potential; Rheo, rheobase; sAHP, slow afterhyperpolarization; AP_{hw}, Action Potential half-width; AP_{amp}, Action Potential amplitude. * $p < 0.05$, ** $p < 0.001$ as compared to pre-E₂ application.

E₂ does not facilitate consolidation of extinction of a cocaine CPP

Previous work demonstrated that E₂ is necessary for the extinction of a CPP (Twining et al., 2013), but did not dissociate the role of E₂ in acquisition versus consolidation of extinction. To dissociate these processes, we focused on the consolidation period immediately after an extinction trial. OVX female rats received systemic injections of either 0.2 mg/kg E₂ (n = 6; Gresack & Frick, 2006) or HBC (n = 7) vehicle immediately after each extinction session (Figure 8). A mixed model three-way ANOVA revealed a significant effect of chamber overall ($F_{(2, 22)} = 10.4, p < 0.01$). However, there was no significant group by chamber by trial interaction ($F_{(8, 88)} = 1.152, p > 0.05$) or trial by chamber effect ($F_{(8, 88)} = 1.623, p > 0.05$). *Post hoc* analysis confirmed that the both E₂-treated and vehicle-treated rats showed a significant preference for the cocaine-paired chamber on trials 1 and 2 (*post hoc* $p < 0.05$). Vehicle-treated rats also showed a significant preference on trial 4 (*post hoc* $p < 0.05$). Thus, post-injections of E₂ did not enhance consolidation of extinction of a cocaine CPP.

Figure 8




Figure 8. Post-injections of E₂ did not facilitate consolidation of extinction of a cocaine CPP. Systemic injections (arrows) of E₂ (n = 6) or vehicle (n = 7) immediately after a CPP trial did not facilitate consolidation of extinction. * $p < 0.05$.

DISCUSSION

The aim of these experiments was to determine the neurotrophic mechanism underlying E₂-induced facilitation of extinction of a cocaine-induced CPP. Based on previous data from the lab (Twining et al., 2013) we expected that E₂, when administered daily, would mediate extinction of cocaine seeking in female rats. Moreover, we hypothesized that E₂-induced facilitation is likely mediated via BDNF/TrkB signaling. However, systemic injections of E₂ prior to each CPP trial did not consistently facilitate extinction (Figure 1). Inconsistent with previous findings (Twining et al., 2013), extinction was evident in the absence of E₂ (Figure 3). Contrary to expectations, direct infusions of the non-selective Trk receptor antagonist, K-252a in IL-mPFC prior to 30 min extinction trials did not impair but facilitated extinction (Figure 1, Figure 2). These results suggest that Trk receptors in IL-mPFC may not play a direct role in mediating the effects of E₂-induced facilitation of extinction. To examine whether TrkB receptors mediated extinction, a highly selective TrkB receptor antagonist, ANA-12 was systemically administered to female rats in conjunction with E₂ (Figure 4). To reduce the variability observed with 30 min extinction trials, 15 min shortened trials were used (Twining et al., 2013). Female rats that received E₂ and vehicle appeared to show facilitated extinction and this effect was not consistently reported in this study. In contrast, rats that were administered E₂ and ANA-12 had impaired extinction of cocaine seeking (Figure 4). The effects of E₂ on the consolidation of extinction of a cocaine CPP were also examined. Post-extinction injections of E₂ did not enhance the consolidation of extinction of a CPP (Figure 8).

Next, we hypothesized that intrinsic excitability would be potentiated by E₂ in IL-mPFC pyramidal neurons and blockade of Trk receptors would attenuate the effects of E₂. OVX female rats received systemic injections of E₂ or HBC vehicle injections prior to being euthanized for

recordings to investigate whether E₂ priming affects membrane properties (Schiess et al., 1988). Bath-application of E₂ enhanced intrinsic excitability in IL-mPFC neurons, independent of E₂ priming (Figure 5, Figure 6). However, this potentiation was significantly attenuated in the presence of the Trk receptor antagonist, K-252a (Figure 7). Overall, we were unable to consistently replicate the original findings of Twining et al., (2013) and show that E₂ is necessary for the acquisition or consolidation of extinction of cocaine seeking. This study suggests that E₂ may be a modulator and not necessarily a mediator of extinction and future studies are required to elucidate the underlying BDNF/TrkB mechanism. However, we were able to support the hypothesis that E₂ enhances intrinsic excitability via BDNF/TrkB signaling in IL-mPFC neurons. This provides evidence that E₂ and TrkB signaling may be modulating neuronal excitability, thereby, facilitating future synaptic plasticity and learning.

Role of Trk receptors in extinction of cocaine seeking

The results of this study were unable to consistently replicate the findings that E₂ is necessary for the acquisition of extinction of a cocaine CPP (Twining et al., 2013). For example, in Figure 1, rats that received IL-mPFC infusions of vehicle and systemic injections of E₂ prior to a 30 min CPP trial showed a significant preference for the cocaine-paired chamber till extinction day 6. A similar trend was also observed in Figure 2 as rats showed a preference for the cocaine-paired chamber till extinction day 4. This data may suggest that E₂ injections before an extinction trial may be necessary to retrieve the original drug memory. This effect has also been observed in a previous study which demonstrated that following conditioning, OVX rats showed significantly low preferences to the cocaine-paired chamber compared to cycling females (Russo et al., 2003). The retrieval deficit was rescued when OVX females were treated with E₂ (Russo et al., 2003).

Thus, these findings and our results suggest that E₂ may play a role in retrieval of the original drug memory rather than enhancing acquisition of extinction.

E₂-facilitated extinction of a cocaine CPP was observed in Figure 4 when rats were exposed to short 15 min extinction trials, instead of 30 min extinction trials. However, when OVX rats were administered E₂ immediately after a 15 min extinction trial, E₂ did not significantly enhance consolidation of extinction compared to rats that received HBC vehicle (Figure 8). Findings from Twining et al., (2013) showed that 15 min extinction trials were sufficient to consistently facilitate extinction induced by E₂. It is possible that the variable effects of E₂ may be due to small sample sizes of 6-8 rats per group. In summary, the present study shows that 30 min extinction trials appear to enhance retrieval of the original drug memory and 15 min extinction trials provide variable results regarding the role of E₂ in extinction. Thus, these experiments need to be repeated with 15 min extinction trials and larger sample sizes.

Inconsistent with previous findings (Twining et al., 2013), extinction was observed in the absence of E₂ (Figure 3). Data from Twining et al., (2013) demonstrated that E₂ is necessary for extinction of a cocaine CPP as E₂-treated rats extinguished within a week, whereas vehicle-treated rats continued to express a CPP for over 5 weeks. In the present study, rats that received IL-mPFC infusions of vehicle and systemic injections of HBC vehicle expressed a CPP on extinction day 1 but did not show preference for the cocaine-paired chamber during subsequent trials (Figure 3). Issues with replication may be due, in part, to the shorter extinction protocols used in the present experiments. The extinction trial protocol used in the previous study (Twining et al., 2013) lasted between 16-32 days in which CPP expression was initially poor but stabilized across days. Thus, future studies require optimizing a reliable short protocol to examine CPP in females to limit variability across studies.

The results of the study did not support our hypothesis that direct Trk receptor blockade in IL-mPFC impairs E₂-facilitated extinction of cocaine seeking. IL-mPFC infusions of a non-selective Trk receptor antagonist, K-252a did not impair but facilitated extinction of a cocaine CPP (Figure 2, Figure 3). A possible explanation for these results may be that K-252a did not affect the new extinction memory, but instead affected the original drug-memory by disrupting reconsolidation. Reconsolidation involves re-stabilizing a memory after the memory has been retrieved and made labile again. For example, following reactivation of an alcohol self-administration memory, administration of NMDA receptor antagonist MK-801 resulted in a strong trend for reconsolidation disruption when tested the following day (Wouda et al., 2010). Similar effects were observed in studies in our lab that aimed to disrupt extinction of a drug-memory in a self-administration model via NMDA receptor antagonists. Infusions of NMDA receptor antagonists appeared to facilitate extinction rather than disrupting it. However, IL-mPFC infusions of NMDA receptor antagonists immediately after short reactivation sessions confirmed that reconsolidation of the original drug memory had been disrupted instead of extinction memory (Hafenbreidel et al., *unpublished*). Thus, even though Trk receptor blockade has previously been shown to disrupt extinction (Xin et al., 2014), these studies indicate that Trk receptors may also be implicated in reconsolidation of the drug-memory.

The present study was partially able to support the hypothesis that TrkB receptor blockade impairs E₂-induced facilitation of extinction. Rats that were administered systemic injections of TrkB receptor antagonist, ANA-12 and E₂ continued to show a preference for the cocaine-paired chamber during short 15 min extinction trials, demonstrating impaired extinction (Figure 4). Rats that received vehicle and E₂ treatment only expressed a CPP on extinction day 1 and continued to show similar preferences for the cocaine- and saline-paired chambers, indicating facilitated

extinction. The results of this particular experiment is in agreement with our hypothesis that TrkB receptor blockade impairs E₂-induced facilitation of extinction. However, it is important to note that we did not observe clear E₂-facilitated extinction in other experiments in this study. To reduce variability in our results, it is necessary to repeat the experiment with larger sample sizes and 15 min extinction trials. Furthermore, to ensure that TrkB receptor blockade impairs extinction, future work should also control for hormonal manipulations and include groups that do not receive E₂ treatment during extinction.

Another possible explanation for the discrepancy in our behavioral data may be stress, which can alter BDNF protein or mRNA expression (Marmigere, Givalois, Rage, Arancibia, & Tapia-Arancibia, 2003; Franklin & Perrot-Sinal, 2006); therefore, E₂ withdrawal or administration may have variable effects on behavior if the rodent has not been handled carefully. Furthermore, prior behavioral handling has also been shown to modify the effects of E₂, and thus, influence changes in spine synapses. For example, OVX rats treated with estradiol benzoate, which also had a 1-day Morris water maze testing did not have increased CA1 spine synapse densities. However, increases in spine synapse densities were observed in behaviorally naïve OVX rats that received hormone treatment (Frick et al., 2004). Thus, stress or prior experiences such as conditioning may have altered the effects of E₂ or BDNF, which may add to the discrepancies in our behavioral data.

A limitation of the present study was that we were unable to test different doses of K-252a or ANA-12 that may impair extinction. Since we did not see differences between groups that were treated with Trk receptor antagonists (Figure 3), we can speculate that the selected doses may not be sufficient to impair memory. The dose for K-252a was chosen from a previous study (Xin et al., 2014), which was infused in IL-mPFC to impair extinction of a conditioned taste aversion

paradigm in male rats. Furthermore, the dose for ANA-12 has previously been used to reduce self-administration of cocaine in male rats (Vassoler, White, Schmidt, Sadri-Vakili, & Pierce, 2013). Thus, due to sex differences or experimental parameters these doses may not apply to impair extinction of a cocaine CPP in female rats. For future studies, it may be important to test different doses of K-252a and ANA-12 on extinction without E₂ administration. Furthermore, the lowest dose of K-252a and ANA-12 that impairs extinction should be chosen to ensure that Trk receptor antagonists do not impair extinction memory on its own but through an E₂-mediated mechanism (Floresco & Jentsch, 2011).

In summary, due to variability in our results, we are unable to conclude that E₂ mediates extinction of cocaine seeking as previously demonstrated by Twining et al., (2013). Issues with replication may be due, in part, to the shorter extinction protocols as we did not run the 16-32 extinction trial protocol used previously. Low sample sizes may also contribute to variability in our data. However, the study was able to demonstrate an interaction between E₂ and BDNF in a CPP paradigm, and this interaction has not been observed in other behavioral paradigms. Thus, the present study provides insight into the E₂-BDNF interaction and its role in learning and memory.

E₂ potentiates intrinsic excitability in IL-mPFC pyramidal neurons via a BDNF-dependent mechanism

Our findings support the hypothesis that E₂ enhances intrinsic excitability in IL-mPFC neurons via a BDNF-dependent mechanism. These findings in IL-mPFC are novel and the first to highlight the E₂ potentiated mechanism underlying extinction of drug seeking or fear conditioning. This study aimed to measure intrinsic excitability which lowers the threshold for synaptic changes and is a neural correlate of learning. A significant amount of attention has been

given to synaptic plasticity as a mechanism for learning. However, evidence from recent studies show that learning and patterns of neuronal activity produce diverse and global changes by regulating membrane properties. Furthermore, changes in excitability are often expressed as changes in resting and voltage-dependent channels and ion pumps (Daoudal & Debanne, 2003; Mozzachiodi & Byrne, 2010; Sehgal, Song, Ehlers, & Moyer, 2013). Thus, enhanced intrinsic excitability has been shown to be correlated with enhanced learning. For example, increased intrinsic excitability has been observed following eyeblink conditioning and trace fear conditioning, (Kaczorowski & Disterhoft, 2009; McKay, Matthews, Oliveira, & Disterhoft, 2009; Song et al., 2012), which are widely used paradigms to study basic mechanisms of learning and memory. Specifically, intrinsic plasticity has been shown to be important for extinction. *In vitro* studies have demonstrated that fear conditioning depressed IL-mPFC excitability, which was demonstrated by a decreased number of APs and increased sAHP. Moreover, extinction reversed the conditioning-induced effects and increased number of APs (Santini, Quirk, & Porter, 2008). Thus, non-synaptic or intrinsic plasticity plays a crucial role in shaping adaptive behaviors such as extinction.

Our results reveal that E₂ enhances intrinsic excitability in IL-mPFC parallel earlier findings from hippocampal pyramidal neurons (Kumar & Foster, 2002; Carrer et al., 2003; Wu et al., 2011), which implicate the crucial role of E₂ in learning and memory. In addition to increasing spontaneous firing in hippocampal neurons, bath-application of E₂ can enhance baseline synaptic responses by increasing the amplitude of excitatory postsynaptic potential (EPSPs) in CA3-CA1 synapses in slices from male and female rodents (Teyler, Vardaris, Lewis, & Rawitch, 1980; Wong & Moss, 1992; Foy et al., 1999; Fugger, Kumar, Lubahn, Korach, & Foster, 2001). E₂ can also initiate a rapid and reversible increase in kainate-induced currents in CA1 hippocampal

neurons (Wong & Moss, 1991, 1992). Enhancement of kainate-induced currents by E₂ was blocked by a blocker of protein kinase A (PKA) suggesting that E₂ may activate intracellular second messenger systems to modulate neuronal function (Gu & Moss, 1996). E₂ has been shown to potentiate long-term potentiation (LTP) in CA3-CA1 synapses in slices from female rats (Bi, Broutman, Foy, Thompson, & Baudry, 2000; Foy, 2001; Woolley, 2007). LTP is the persistent strengthening of synapses and the underlying mechanism of synaptic plasticity, learning, and memory (Malinow, Schulman, & Tsien, 1989). Thus, the short and long-term effects of extinction may be mediated by E₂-enhanced intrinsic excitability and synaptic potentiation.

Similar to the effects of E₂ on synaptic plasticity, perfusion of BDNF induces LTP in the hippocampus (Ying et al., 2002; Ji et al., 2010). The role of Trk receptors in LTP have also been shown in mice carrying a mutated TrkB receptor (TrkB^{2478A/2478A}), which cannot be phosphorylated. Brain slices from TrkB^{2478A/2478A} mice show reductions in late LTP in CA3-CA1 synapses (Lai et al., 2012). However, the role of BDNF or Trk receptors in intrinsic excitability remains unclear. For example, BDNF has shown to reduce excitability in visual cortical cultures (Desai, Rutherford, & Turrigiano, 1999), but TrkB receptor activation enhanced excitability in slices or cortical neurons (Marongiu et al., 2013). TrkB receptor activation also enhanced NMDA-receptor mediated currents in IL-mPFC neurons from male rats (Otis et al., 2014). Our results advance the findings by demonstrating the role of Trk receptors in non-synaptic plasticity in IL-mPFC, a region crucial for extinction.

In summary, our data reveal that E₂ enhances non-synaptic plasticity in IL-mPFC pyramidal neurons. Previous work has been predominantly focused on E₂-enhanced excitability and synaptic activity in the hippocampus (Wong & Moss, 1991; Kumar & Foster, 2002; Carrer et al., 2003). Thus, our findings that E₂-induced potentiation of intrinsic excitability in IL-mPFC

complements previous work in the hippocampus. Since both the hippocampus and IL-mPFC are engaged in extinction, these findings suggest that E₂ plays a critical role in modulating extinction. Finally, our results suggest that E₂-enhanced excitability in IL-mPFC neurons is mediated by BDNF/TrkB receptor activation and this interaction may likely regulate extinction across paradigms.

Future directions

Despite the currently conflicting data regarding the role of E₂ in extinction of a cocaine CPP, previous research shows that E₂ enhances performance on a wide range of cognitive tasks as well as extinction in other paradigms (Luine et al., 2003; Milad et al., 2009; Tuscher et al., 2015). E₂ replacement or elevated E₂ levels are associated with improved spatial reference memory in a water maze task (Gresack & Frick, 2006), object recognition (Lewis et al., 2008; Jacome et al., 2010), and extinction of conditioned fear in female rats and humans (Zeidan et al., 2011; Graham & Milad, 2013). Conversely, low levels of E₂ are associated with extinction failure in females with post-traumatic stress disorder (Glover et al., 2012). Furthermore, low levels of E₂ in both rats and humans leads to extinction impairment of conditioned fear (Graham & Milad, 2013). Thus, the effects of E₂-induced enhancement of extinction that vary in a CPP paradigm, may be more robust in other paradigms such as fear conditioning, or cocaine self-administration

In addition to investigating the role of BDNF as an underlying mechanism for E₂-induced facilitation of extinction, it is important to target another neurotrophic factor, basic fibroblast growth factor (bFGF). Studies conducted in our lab have shown that cocaine-related learning increases bFGF protein expression in IL-mPFC. Furthermore, blocking bFGF in IL-mPFC reverses this enhanced expression and facilitates extinction of cocaine seeking (Hafenbreidel, Twining, Rafa Todd, & Mueller, 2015). bFGF has also shown to reduce intrinsic excitability in

hippocampal pyramidal neurons by inhibiting voltage-gated Na⁺ (Hilborn, Vaillancourt, Rane, 1998) and K⁺ currents (Cuppini, Ambrogini, Lattanzi, Ciuffoli, & Cuppini, 2009). Reduced excitability is associated with decreased synaptic plasticity (Gelinias & Nguyen, 2005), thus, increased bFGF expression would potentially reduce extinction-related synaptic plasticity. Previously, E₂ has been shown to interact with bFGF as loss of circulating E₂ resulted in high bFGF expression (Flores, Salmaso, Cain, Rodaros, & Stewart, 1999). E₂ replacement reduced bFGF expression (Flores et al., 1999), which suggests that E₂ may enhance extinction by reversing the maladaptive overexpression of E₂ observed with cocaine-related learning. E₂ may also have the potential to reduce intrinsic excitability and promote extinction-related plasticity. Thus, bFGF may be a promising target to elucidate the mnemonic and neural mechanisms, underlying E₂-induced enhancement of extinction.

CONCLUSION

Persistent drug seeking is maintained by drug-associated memories, through which cues can elicit craving and relapse. Impairing drug-associated memories may dampen the motivation to seek drugs and limit relapse susceptibility. Specifically, women are more vulnerable than males to key phases of addiction, which include initiation, bingeing, and relapse (Anker & Carroll, 2010). Rehabilitation programs that use exposure therapy, in which addicts are exposed to a drug-associated cue in the absence of drug reinforcement have had limited success without any pharmacological adjuncts (Conklin & Tiffany, 2002). Thus, pharmacological enhancement of E₂ or BDNF/TrkB signaling may prove to be clinically relevant for treatment of disorders involving maladaptive memories and behavioral inflexibility such as addiction or posttraumatic stress disorder.

REFERENCES

- Acosta, J. I., Mayer, L., Talboom, J. S., Tsang, C. W. S., Smith, C. J., Enders, C. K., & Bimonte-Nelson, H. A. (2009). Transitional versus surgical menopause in a rodent model: etiology of ovarian hormone loss impacts memory and the acetylcholine system. *Endocrinology*, *150*(9), 4248-4259.
- Aguirre, C. C., & Baudry, M. (2009). Progesterone reverses 17 β -estradiol-mediated neuroprotection and BDNF induction in cultured hippocampal slices. *European Journal of Neuroscience*, *29*(3), 447-454.
- Allen, A. L., & McCarson, K. E. (2005). Estrogen increases nociception-evoked brain-derived neurotrophic factor gene expression in the female rat. *Neuroendocrinology*, *81*(3), 193-199.
- Almey, A., Cannell, E., Bertram, K., Filardo, E., Milner, T. A., & Brake, W. G. (2014). Medial prefrontal cortical estradiol rapidly alters memory system bias in female rats: ultrastructural analysis reveals membrane-associated estrogen receptors as potential mediators. *Endocrinology*, *155*(11), 4422-4432.
- Anker, J. J., & Carroll, M. E. (2010). Females are more vulnerable to drug abuse than males: evidence from preclinical studies and the role of ovarian hormones. In *Biological basis of sex differences in psychopharmacology* (pp. 73-96). Springer Berlin Heidelberg.
- Back, S. E., Brady, K. T., Jackson, J. L., Salstrom, S., & Zinzow, H. (2005). Gender differences in stress reactivity among cocaine-dependent individuals. *Psychopharmacology*, *180*(1), 169-176.
- Becker, J. B., & Hu, M. (2008). Sex differences in drug abuse. *Frontiers in neuroendocrinology*, *29*(1), 36-47.
- Berglind, W. J., See, R. E., Fuchs, R. A., Ghee, S. M., Whitfield, T. W., Miller, S. W., & McGinty, J. F. (2007). A BDNF infusion into the medial prefrontal cortex suppresses cocaine seeking in rats. *European Journal of Neuroscience*, *26*(3), 757-766.
- Bi, R., Broutman, G., Foy, M. R., Thompson, R. F., & Baudry, M. (2000). The tyrosine kinase and mitogen-activated protein kinase pathways mediate multiple effects of estrogen in hippocampus. *Proceedings of the National Academy of Sciences*, *97*(7), 3602-3607.
- Bian, J. T., Zhang, J. W., Zhang, Z. X., & Zhao, H. L. (2005). Association analysis of brain-derived neurotrophic factor (BDNF) gene 196 A/G polymorphism with Alzheimer's disease (AD) in mainland Chinese. *Neuroscience letters*, *387*(1), 11-16.
- Bimonte, H. A., & Denenberg, V. H. (1999). Estradiol facilitates performance as working memory load increases. *Psychoneuroendocrinology*, *24*(2), 161-173.
- Bimonte-Nelson, H. A., Nelson, M. E., & Granholm, A. C. E. (2004). Progesterone counteracts estrogen-induced increases in neurotrophins in the aged female rat brain. *Neuroreport*, *15*(17), 2659-2663.

- Blanquet, P. R., Mariani, J., & Derer, P. (2003). A calcium/calmodulin kinase pathway connects brain-derived neurotrophic factor to the cyclic AMP-responsive transcription factor in the rat hippocampus. *Neuroscience*, *118*(2), 477-490.
- Bobzean, S. A., Dennis, T. S., & Perrotti, L. I. (2014). Acute estradiol treatment affects the expression of cocaine-induced conditioned place preference in ovariectomized female rats. *Brain research bulletin*, *103*, 49-53.
- Boulware, M. I., Weick, J. P., Becklund, B. R., Kuo, S. P., Groth, R. D., & Mermelstein, P. G. (2005). Estradiol activates group I and II metabotropic glutamate receptor signaling, leading to opposing influences on cAMP response element-binding protein. *The Journal of neuroscience*, *25*(20), 5066-5078.
- Boulware, M. I., Heisler, J. D., & Frick, K. M. (2013). The memory-enhancing effects of hippocampal estrogen receptor activation involve metabotropic glutamate receptor signaling. *The Journal of Neuroscience*, *33*(38), 15184-15194.
- Bourne, J. N., & Harris, K. M. (2008). Balancing structure and function at hippocampal dendritic spines. *Annual review of neuroscience*, *31*, 47.
- Bouton ME, Garcia-Gutierrez A, Zilski J, Moody EW (2006) Extinction in multiple contexts does not necessarily make extinction less vulnerable to relapse. *Behaviour Research and Therapy* *44*, 983–994.
- Bouton, M. E., & Todd, T. P. (2014). A fundamental role for context in instrumental learning and extinction. *Behavioural processes*, *104*, 13-19.
- Brady, K. T., & Randall, C. L. (1999). Gender differences in substance use disorders. *Psychiatric Clinics of North America*, *22*(2), 241-252.
- Brann, D. W., Dhandapani, K., Wakade, C., Mahesh, V. B., & Khan, M. M. (2007). Neurotrophic and neuroprotective actions of estrogen: basic mechanisms and clinical implications. *Steroids*, *72*(5), 381-405.
- Carrer, H. F., Araque, A., & Buño, W. (2003). Estradiol regulates the slow Ca²⁺-activated K⁺ current in hippocampal pyramidal neurons. *The Journal of neuroscience*, *23*(15), 6338-6344.
- Carroll, M. E., Lynch, W. J., Roth, M. E., Morgan, A. D., & Cosgrove, K. P. (2004). Sex and estrogen influence drug abuse. *Trends in pharmacological sciences*, *25*(5), 273-279.
- Carter, B. L., & Tiffany, S. T. (1999). Meta-analysis of cue-reactivity in addiction research. *Addiction*, *94*(3), 327-340.
- Cazorla, M., Prémont, J., Mann, A., Girard, N., Kellendonk, C., & Rognan, D. (2011). Identification of a low-molecular weight TrkB antagonist with anxiolytic and antidepressant activity in mice. *The Journal of clinical investigation*, *121*(5), 1846-1857.

- Chao, M. V. (2003). Neurotrophins and their receptors: a convergence point for many signalling pathways. *Nature Reviews Neuroscience*, 4(4), 299-309.
- Chen, K., & Kandel, D. (2002). Relationship between extent of cocaine use and dependence among adolescents and adults in the United States. *Drug and Alcohol Dependence*, 68(1), 65-85.
- Chesworth, R., and Corbit, L. H. (2015). Recent developments in the behavioural and pharmacological enhancement of extinction of drug seeking. *Addiction Biology*, doi: [10.1111/adb.12337](https://doi.org/10.1111/adb.12337).
- Christensen, A., Dewing, P., & Micevych, P. (2011). Membrane-initiated estradiol signaling induces spinogenesis required for female sexual receptivity. *The Journal of Neuroscience*, 31(48), 17583-17589.
- Clipperton-Allen, A. E., Almey, A., Melicherik, A., Allen, C. P., & Choleris, E. (2011). Effects of an estrogen receptor alpha agonist on agonistic behaviour in intact and gonadectomized male and female mice. *Psychoneuroendocrinology*, 36(7), 981-995.
- Cohen, S. J., Munchow, A. H., Rios, L. M., Zhang, G., Ásgeirsdóttir, H. N., & Stackman, R. W. (2013). The rodent hippocampus is essential for nonspatial object memory. *Current Biology*, 23(17), 1685-1690.
- Cohen-Cory, S., Kidane, A. H., Shirkey, N. J., & Marshak, S. (2010). Brain-derived neurotrophic factor and the development of structural neuronal connectivity. *Developmental neurobiology*, 70(5), 271-288.
- Conklin, C. A., & Tiffany, S. T. (2002). Applying extinction research and theory to cue-exposure addiction treatments. *Addiction*, 97(2), 155-167.
- Corcoran, K. A., & Maren, S. (2001). Hippocampal inactivation disrupts contextual retrieval of fear memory after extinction. *The Journal of neuroscience*, 21(5), 1720-1726.
- Cornil, C. A., & Charlier, T. D. (2010). Rapid behavioural effects of oestrogens and fast regulation of their local synthesis by brain aromatase. *Journal of neuroendocrinology*, 22(7), 664-673.
- Cuppini, C., Ambrogini, P., Lattanzi, D., Ciuffoli, S., & Cuppini, R. (2009). FGF2 modulates the voltage-dependent K⁺ current and changes excitability of rat dentate gyrus granule cells. *Neuroscience letters*, 462(3), 203-206.
- Daniel, J. M., Fader, A. J., Spencer, A. L., & Dohanich, G. P. (1997). Estrogen enhances performance of female rats during acquisition of a radial arm maze. *Hormones and behavior*, 32(3), 217-225.
- Daniel, J. M., Roberts, S. L., & Dohanich, G. P. (1999). Effects of ovarian hormones and environment on radial maze and water maze performance of female rats. *Physiology & Behavior*, 66(1), 11-20.

- Daoudal, G., & Debanne, D. (2003). Long-term plasticity of intrinsic excitability: learning rules and mechanisms. *Learning & Memory*, 10(6), 456-465.
- Davis, W. M., & Smith, S. G. (1977). Catecholaminergic mechanisms of reinforcement: Direct assessment by drug self-administration. *Life sciences*, 20(3), 483-492.
- Desai, N. S., Rutherford, L. C., & Turrigiano, G. G. (1999). BDNF regulates the intrinsic excitability of cortical neurons. *Learning & Memory*, 6(3), 284-291.
- Dohanich, G. P. (2002). Gonadal steroids, learning and memory. *Hormones, brain and behavior*, 2, 265-327.
- Eisenberg, M., & Dudai, Y. (2004). Reconsolidation of fresh, remote, and extinguished fear memory in Medaka: old fears don't die. *European Journal of Neuroscience*, 20(12), 3397-3403.
- Elman, I., Karlsgodt, K. H., & Gastfriend, D. R. (2001). Gender differences in cocaine craving among non-treatment-seeking individuals with cocaine dependence. *The American journal of drug and alcohol abuse*, 27(2), 193-202.
- Ernfors, P., & Bramham, C. R. (2003). The coupling of a trkB tyrosine residue to LTP. *Trends in neurosciences*, 26(4), 171-173.
- Fader, A. J., Johnson, P. E., & Dohanich, G. P. (1999). Estrogen improves working but not reference memory and prevents amnesic effects of scopolamine on a radial-arm maze. *Pharmacology Biochemistry and Behavior*, 62(4), 711-717.
- Fernandez, S. M., Lewis, M. C., Pechenino, A. S., Harburger, L. L., Orr, P. T., Gresack, J. E., ... & Frick, K. M. (2008). Estradiol-induced enhancement of object memory consolidation involves hippocampal extracellular signal-regulated kinase activation and membrane-bound estrogen receptors. *The Journal of Neuroscience*, 28(35), 8660-8667.
- Flores, C., Salmaso, N., Cain, S., Rodaros, D., & Stewart, J. (1999). Ovariectomy of adult rats leads to increased expression of astrocytic basic fibroblast growth factor in the ventral tegmental area and in dopaminergic projection regions of the entorhinal and prefrontal cortex. *The Journal of neuroscience*, 19(19), 8665-8673.
- Floresco, S. B., & Jentsch, J. D. (2011). Pharmacological enhancement of memory and executive functioning in laboratory animals. *Neuropsychopharmacology*, 36(1), 227-250.
- Fonseca, C. S., Gusmão, I. D., Raslan, A. C., Monteiro, B. M. M., Massensini, A. R., Moraes, M.F., & Pereira, G. S. (2013). Object recognition memory and temporal lobe activation after delayed estrogen replacement therapy. *Neurobiology of learning and memory*, 101, 19-25.
- Fortress, A. M., Fan, L., Orr, P. T., Zhao, Z., & Frick, K. M. (2013). Estradiol-induced object recognition memory consolidation is dependent on activation of mTOR signaling in the dorsal hippocampus. *Learning & memory*, 20(3), 147-155.

- Fortress, A. M., Kim, J., Poole, R. L., Gould, T. J., & Frick, K. M. (2014). 17β -Estradiol regulates histone alterations associated with memory consolidation and increases Bdnf promoter acetylation in middle-aged female mice. *Learning & Memory*, *21*(9), 457-467.
- Foy, M. R., Xu, J., Xie, X., Brinton, R. D., Thompson, R. F., & Berger, T. W. (1999). 17β -estradiol enhances NMDA receptor-mediated EPSPs and long-term potentiation. *Journal of Neurophysiology*, *81*(2), 925-929.
- Foy, M. R. (2001). 17β -estradiol: effect on CA1 hippocampal synaptic plasticity. *Neurobiology of learning and memory*, *76*(3), 239-252.
- Franken, I. H., de Haan, H. A., van der Meer, C. W., Haffmans, P. J., & Hendriks, V. M. (1999). Cue reactivity and effects of cue exposure in abstinent posttreatment drug users. *Journal of Substance Abuse Treatment*, *16*(1), 81-85.
- Franklin, T. B., & Perrot-Sinal, T. S. (2006). Sex and ovarian steroids modulate brain-derived neurotrophic factor (BDNF) protein levels in rat hippocampus under stressful and non-stressful conditions. *Psychoneuroendocrinology*, *31*(1), 38-48.
- Frick, K. M., Fernandez, S. M., Bennett, J. C., Prange-Kiel, J., MacLusky, N. J., & Leranth, C. (2004). Behavioral training interferes with the ability of gonadal hormones to increase CA1 spine synapse density in ovariectomized female rats. *European Journal of neuroscience*, *19*(11), 3026-3032.
- Frick, A., & Johnston, D. (2005). Plasticity of dendritic excitability. *Journal of neurobiology*, *64*(1), 100-115.
- Frick, K. M. (2009). Estrogens and age-related memory decline in rodents: what have we learned and where do we go from here?. *Hormones and behavior*, *55*(1), 2-23.
- Frick, K. M., Fernandez, S. M., & Harburger, L. L. (2010). A new approach to understanding the molecular mechanisms through which estrogens affect cognition. *Biochimica et Biophysica Acta (BBA)-General Subjects*, *1800*(10), 1045-1055.
- Frick, K. M. (2012). Building a better hormone therapy? How understanding the rapid effects of sex steroid hormones could lead to new therapeutics for age-related memory decline. *Behavioral neuroscience*, *126*(1), 29.
- Frick, K. M. (2015). Molecular mechanisms underlying the memory-enhancing effects of estradiol. *Hormones and behavior*, *74*, 4-18.
- Frick, K. M., Kim, J., Tuscher, J. J., & Fortress, A. M. (2015). Sex steroid hormones matter for learning and memory: estrogenic regulation of hippocampal function in male and female rodents. *Learn Mem*, *22*(9), 472-93.
- Fugger, H. N., Kumar, A., Lubahn, D. B., Korach, K. S., & Foster, T. C. (2001). Examination of estradiol effects on the rapid estradiol mediated increase in hippocampal synaptic transmission in estrogen receptor α knockout mice. *Neuroscience letters*, *309*(3), 207-209.

- Garza-Meilandt, A., Cantu, R. E., & Claiborne, B. J. (2006). Estradiol's effects on learning and neuronal morphology vary with route of administration. *Behavioral neuroscience*, *120*(4), 905-916.
- Gelinas, J. N., & Nguyen, P. V. (2005). β -Adrenergic receptor activation facilitates induction of a protein synthesis-dependent late phase of long-term potentiation. *The Journal of neuroscience*, *25*(13), 3294-3303.
- Gibbs, R. B. (1998). Levels of trkA and BDNF mRNA, but not NGF mRNA, fluctuate across the estrous cycle and increase in response to acute hormone replacement. *Brain research*, *787*(2), 259-268.
- Gibbs, R. B. (1999). Treatment with estrogen and progesterone affects relative levels of brain-derived neurotrophic factor mRNA and protein in different regions of the adult rat brain. *Brain research*, *844*(1), 20-27.
- Gibbs, R. B., & Johnson, D. A. (2008). Sex-specific effects of gonadectomy and hormone treatment on acquisition of a 12-arm radial maze task by Sprague Dawley rats. *Endocrinology*, *149*(6), 3176-3183.
- Gillies, G. E., & McArthur, S. (2010). Estrogen actions in the brain and the basis for differential action in men and women: a case for sex-specific medicines. *Pharmacological reviews*, *62*(2), 155-198.
- Glover, E. M., Jovanovic, T., Mercer, K. B., Kerley, K., Bradley, B., Ressler, K. J., & Norrholm, S. D. (2012). Estrogen levels are associated with extinction deficits in women with posttraumatic stress disorder. *Biological Psychiatry*, *72*(1), 19-24.
- Gottmann, K., Mittmann, T., & Lessmann, V. (2009). BDNF signaling in the formation, maturation and plasticity of glutamatergic and GABAergic synapses. *Experimental brain research*, *199*(3-4), 203-234.
- Gouveia, A., dos Santos, U. D., Felisbino, F. E., de Afonseca, T. L., Antunes, G., & Morato, S. (2004). Influence of the estrous cycle on the behavior of rats in the elevated T-maze. *Behavioural processes*, *67*(2), 167-171.
- Graham, B. M., & Milad, M. R. (2013). Blockade of estrogen by hormonal contraceptives impairs fear extinction in female rats and women. *Biological psychiatry*, *73*(4), 371-378.
- Graham, B. M., & Milad, M. R. (2014). Inhibition of estradiol synthesis impairs fear extinction in male rats. *Learning & Memory*, *21*(7), 347-350.
- Gresack, J. E., & Frick, K. M. (2004). Environmental enrichment reduces the mnemonic and neural benefits of estrogen. *Neuroscience*, *128*(3), 459-471.
- Gresack, J. E., & Frick, K. M. (2006). Post-training estrogen enhances spatial and object memory consolidation in female mice. *Pharmacology Biochemistry and Behavior*, *84*(1), 112-119.

- Gu, Q., & Moss, R. L. (1996). 17 β -Estradiol potentiates kainate-induced currents via activation of the cAMP cascade. *The Journal of neuroscience*, 16(11), 3620-3629.
- Hafenbreidel, M., Todd, C. R., Twining, R. C., Tuscher, J. J., & Mueller, D. (2014). Bidirectional effects of inhibiting or potentiating NMDA receptors on extinction after cocaine self-administration in rats. *Psychopharmacology*, 231(24), 4585-4594.
- Hafenbreidel, M., Twining, R. C., Todd, C. R., & Mueller, D. (2015). Blocking infralimbic basic fibroblast growth factor (bFGF or FGF2) facilitates extinction of drug seeking after cocaine self-administration. *Neuropsychopharmacology*, 40(13), 2907-2915.
- Hamilton, R. T., Rettberg, J. R., Mao, Z., To, J., Zhao, L., Appt, S. E., ... & Brinton, R. D. (2011). Hippocampal responsiveness to 17 β -estradiol and equol after long-term ovariectomy: Implication for a therapeutic window of opportunity. *Brain research*, 1379, 11-22.
- Heikkinen, T., Puoliväli, J., Liu, L., Rissanen, A., & Tanila, H. (2002). Effects of ovariectomy and estrogen treatment on learning and hippocampal neurotransmitters in mice. *Hormones and behavior*, 41(1), 22-32.
- Hill, R. A. (2012). Interaction of Sex Steroid Hormones and Brain-Derived Neurotrophic Factor-Tyrosine Kinase B Signalling: Relevance to Schizophrenia and Depression. *Journal of neuroendocrinology*, 24(12), 1553-1561.
- Hilborn, M. D., Vaillancourt, R. R., & Rane, S. G. (1998). Growth factor receptor tyrosine kinases acutely regulate neuronal sodium channels through the src signaling pathway. *The Journal of neuroscience*, 18(2), 590-600.
- Holmes, M. M., Wide, J. K., & Galea, L. A. (2002). Low levels of estradiol facilitate, whereas high levels of estradiol impair, working memory performance on the radial arm maze. *Behavioral Neuroscience*, 116(5), 928.
- Hu, M., & Becker, J. B. (2003). Effects of sex and estrogen on behavioral sensitization to cocaine in rats. *The Journal of neuroscience*, 23(2), 693-699.
- Hu, M., Crombag, H. S., Robinson, T. E., & Becker, J. B. (2004). Biological basis of sex differences in the propensity to self-administer cocaine. *Neuropsychopharmacology*, 29(1), 81-85.
- Hu, M., & Becker, J. B. (2008). Acquisition of cocaine self-administration in ovariectomized female rats: effect of estradiol dose or chronic estradiol administration. *Drug and alcohol dependence*, 94(1), 56-62.
- Huang, E. J., & Reichardt, L. F. (2003). Trk receptors: roles in neuronal signal transduction*. *Annual review of biochemistry*, 72(1), 609-642.
- Ignjatova, L., & Raleva, M. (2008). Gender difference in the treatment outcome of patients served in the mixed-gender program. *Bratislavské lekárske listy*, 110(5), 285-289.

- Inagaki, T., Gautreaux, C., & Luine, V. (2010). Acute estrogen treatment facilitates recognition memory consolidation and alters monoamine levels in memory-related brain areas. *Hormones and behavior*, 58(3), 415-426.
- Jackson, L. R., Robinson, T. E., & Becker, J. B. (2006). Sex differences and hormonal influences on acquisition of cocaine self-administration in rats. *Neuropsychopharmacology*, 31(1), 129-138.
- Jacome, L. F., Gautreaux, C., Inagaki, T., Mohan, G., Alves, S., Lubbers, L. S., & Luine, V. (2010). Estradiol and ER β agonists enhance recognition memory, and DPN, an ER β agonist, alters brain monoamines. *Neurobiology of learning and memory*, 94(4), 488-498.
- Jeziarski, M. K., & Sohrabji, F. (2000). Region-and peptide-specific regulation of the neurotrophins by estrogen. *Molecular Brain Research*, 85(1), 77-84.
- Jeziarski, M. K., & Sohrabji, F. (2001). Neurotrophin expression in the reproductively senescent forebrain is refractory to estrogen stimulation. *Neurobiology of aging*, 22(2), 311-321.
- Ji, Y., Lu, Y., Yang, F., Shen, W., Tang, T. T. T., Feng, L., ... & Lu, B. (2010). Acute and gradual increases in BDNF concentration elicit distinct signaling and functions in neurons. *Nature neuroscience*, 13(3), 302-309.
- Jones, M. E., Boon, W. C., Proietto, J., & Simpson, E. R. (2006). Of mice and men: the evolving phenotype of aromatase deficiency. *Trends in Endocrinology & Metabolism*, 17(2), 55-64.
- Kaczorowski, C. C., & Disterhoft, J. F. (2009). Memory deficits are associated with impaired ability to modulate neuronal excitability in middle-aged mice. *Learning & Memory*, 16(6), 362-366.
- Kaczorowski, C. C., Davis, S. J., & Moyer, J. R. (2012). Aging redistributes medial prefrontal neuronal excitability and impedes extinction of trace fear conditioning. *Neurobiology of aging*, 33(8), 1744-1757.
- Kandel, D., Chen, K., Warner, L. A., Kessler, R. C., & Grant, B. (1997). Prevalence and demographic correlates of symptoms of last year dependence on alcohol, nicotine, marijuana and cocaine in the U.S. population. *Drug Alcohol Dependence*, 44(1), 11-29.
- Kerstetter, K. A., Aguilar, V. R., Parrish, A. B., & Kippin, T. E. (2008). Protracted time-dependent increases in cocaine seeking behavior during cocaine withdrawal in female relative to male rats. *Psychopharmacology*, 198(1), 63-75.
- Kippin, T. E., Fuchs, R. A., Mehta, R. H., Case, J. M., Parker, M. P., Bimonte-Nelson, H. A., & See, R. E. (2005). Potentiation of cocaine-primed reinstatement of drug seeking in female rats during estrus. *Psychopharmacology*, 182(2), 245-252.
- Kosten, T. A., Gawin, F. H., Kosten, T. R., & Rounsaville, B. J. (1993). Gender differences in cocaine use and treatment response. *Journal of substance abuse treatment*, 10(1), 63-66.

- Kuiper, G. G., Shughrue, P. J., Merchenthaler, I., & Gustafsson, J. Å. (1998). The estrogen receptor β subtype: a novel mediator of estrogen action in neuroendocrine systems. *Frontiers in neuroendocrinology*, *19*(4), 253-286.
- Kumar, A., & Foster, T. C. (2002). 17β -estradiol benzoate decreases the AHP amplitude in CA1 pyramidal neurons. *Journal of Neurophysiology*, *88*(2), 621-626.
- Lai, K. O., Wong, A. S., Cheung, M. C., Xu, P., Liang, Z., Lok, K. C., ... & Cheung, Z. H. (2012). TrkB phosphorylation by Cdk5 is required for activity-dependent structural plasticity and spatial memory. *Nature neuroscience*, *15*(11), 1506-1515.
- LaLumiere, R. T., Niehoff, K. E., & Kalivas, P. W. (2010). The infralimbic cortex regulates the consolidation of extinction after cocaine self-administration. *Learning & memory*, *17*(4), 168-175.
- Larson, E. B., Anker, J. J., Gliddon, L. A., Fons, K. S., & Carroll, M. E. (2007). Effects of estrogen and progesterone on the escalation of cocaine self-administration in female rats during extended access. *Experimental and clinical psychopharmacology*, *15*(5), 461.
- Lejuez, C. W., Bornovalova, M. A., Reynolds, E. K., Daughters, S. B., & Curtin, J. J. (2007). Risk factors in the relationship between gender and crack/cocaine. *Experimental and clinical psychopharmacology*, *15*(2), 165.
- Levin, E. R. (2005). Integration of the extranuclear and nuclear actions of estrogen. *Molecular Endocrinology*, *19*(8), 1951-1959.
- Lewis, M. C., Kerr, K. M., Orr, P. T., & Frick, K. M. (2008). Estradiol-induced enhancement of object memory consolidation involves NMDA receptors and protein kinase A in the dorsal hippocampus of female C57BL/6 mice. *Behavioral neuroscience*, *122*(3), 716.
- Li, C., Brake, W. G., Romeo, R. D., Dunlop, J. C., Gordon, M., Buzescu, R., ... & McEwen, B. S. (2004). Estrogen alters hippocampal dendritic spine shape and enhances synaptic protein immunoreactivity and spatial memory in female mice. *Proceedings of the National Academy of Sciences*, *101*(7), 2185-2190.
- Likhtik, E., Popa, D., Apergis-Schoute, J., Fidacaro, G. A., & Paré, D. (2008). Amygdala intercalated neurons are required for expression of fear extinction. *Nature*, *454*(7204), 642-645.
- Linnarsson, S., Björklund, A., & Ernfors, P. (1997). Learning deficit in BDNF mutant mice. *European Journal of Neuroscience*, *9*(12), 2581-2587.
- Liu, Y., Fowler, C. D., Young, L. J., Yan, Q., Insel, T. R., & Wang, Z. (2001). Expression and estrogen regulation of brain-derived neurotrophic factor gene and protein in the forebrain of female prairie voles. *Journal of Comparative Neurology*, *433*(4), 499-514.
- Liu, F., Day, M., Muniz, L. C., Bitran, D., Arias, R., Revilla-Sanchez, R., ... & Sung, A. (2008). Activation of estrogen receptor- β regulates hippocampal synaptic plasticity and improves memory. *Nature neuroscience*, *11*(3), 334-343.

- Lowry, N. C., Pardon, L. P., Yates, M. A., & Juraska, J. M. (2010). Effects of long-term treatment with 17 β -estradiol and medroxyprogesterone acetate on water maze performance in middle aged female rats. *Hormones and behavior*, *58*(2), 200-207.
- Lu, B. (2003). BDNF and activity-dependent synaptic modulation. *Learning & memory*, *10*(2), 86-98.
- Luine, V. N., Richards, S. T., Wu, V. Y., & Beck, K. D. (1998). Estradiol enhances learning and memory in a spatial memory task and effects levels of monoaminergic neurotransmitters. *Hormones and behavior*, *34*(2), 149-162.
- Luine, V. N., Jacome, L. F., & MacLusky, N. J. (2003). Rapid enhancement of visual and place memory by estrogens in rats. *Endocrinology*, *144*(7), 2836-2844.
- Luine, V., & Frankfurt, M. (2013). Interactions between estradiol, BDNF and dendritic spines in promoting memory. *Neuroscience*, *239*, 34-45.
- Lynch, W. J., Arizzi, M. N., & Carroll, M. E. (2000). Effects of sex and the estrous cycle on regulation of intravenously self-administered cocaine in rats. *Psychopharmacology*, *152*(2), 132-139.
- Lynch, W. J., Roth, M. E., Mickelberg, J. L., & Carroll, M. E. (2001). Role of estrogen in the acquisition of intravenously self-administered cocaine in female rats. *Pharmacology Biochemistry and Behavior*, *68*(4), 641-646.
- Lynch, W. J., Roth, M. E., & Carroll, M. E. (2002). Biological basis of sex differences in drug abuse: preclinical and clinical studies. *Psychopharmacology*, *164*(2), 121-137.
- Lynch, W. J., & Taylor, J. R. (2004). Sex differences in the behavioral effects of 24-h/day access to cocaine under a discrete trial procedure. *Neuropsychopharmacology*, *29*, 943-951.
- Lynch, W. J., Nicholson, K. L., Dance, M. E., Morgan, R. W., & Foley, P. L. (2010). Animal Models of Substance Abuse and Addiction: Implications for Science, Animal Welfare, and Society. *Comparative Medicine*, *60*(3), 177-188.
- Malinow, R., Schulman, H., & Tsien, R. W. (1989). Inhibition of postsynaptic PKC or CaMKII blocks induction but not expression of LTP. *Science*, *245*(4920), 862-866.
- Marcus, S. M., Young, E. A., Kerber, K. B., Kornstein, S., Farabaugh, A. H., Mitchell, J., ... & Rush, A. J. (2005). Gender differences in depression: findings from the STAR* D study. *Journal of affective disorders*, *87*(2), 141-150.
- Markham, J. A., Pych, J. C., & Juraska, J. M. (2002). Ovarian hormone replacement to aged ovariectomized female rats benefits acquisition of the morris water maze. *Hormones and Behavior*, *42*(3), 284-293.
- Markham, J. A., & Juraska, J. M. (2007). Social recognition memory: influence of age, sex, and ovarian hormonal status. *Physiology & behavior*, *92*(5), 881-888.

- Markowska, A. L., & Savonenko, A. V. (2002). Effectiveness of estrogen replacement in restoration of cognitive function after long-term estrogen withdrawal in aging rats. *The Journal of Neuroscience*, 22(24), 10985-10995.
- Marongiu, D., Imbrosci, B., & Mittmann, T. (2013). Modulatory effects of the novel TrkB receptor agonist 7, 8-dihydroxyflavone on synaptic transmission and intrinsic neuronal excitability in mouse visual cortex in vitro. *European journal of pharmacology*, 709(1), 64-71.
- Marmigère, F., Givalois, L., Rage, F., Arancibia, S., & Tapia-Arancibia, L. (2003). Rapid induction of BDNF expression in the hippocampus during immobilization stress challenge in adult rats. *Hippocampus*, 13(5), 646-655.
- Matsushita, S., Arai, H., Matsui, T., Yuzuriha, T., Urakami, K., Masaki, T., & Higuchi, S. (2005). Brain-derived neurotrophic factor gene polymorphisms and Alzheimer's disease. *Journal of neural transmission*, 112(5), 703-771.
- McCance-Katz, E. F., Carroll, K. M., & Rounsaville, B. J. (1999). Gender differences in treatment-seeking cocaine abusers—implications for treatment and prognosis. *The American Journal on Addictions*, 8(4), 300-311.
- McEwen, B. S., & Alves, S. E. (1999). Estrogen actions in the central nervous system 1. *Endocrine reviews*, 20(3), 279-307.
- McEwen, B., Akama, K., Alves, S., Brake, W. G., Bulloch, K., Lee, S., ... & Milner, T. A. (2001). Tracking the estrogen receptor in neurons: implications for estrogen-induced synapse formation. *Proceedings of the National Academy of Sciences*, 98(13), 7093-7100.
- McKay, J. R., Rutherford, M. J., Cacciola, J. S., Kabasakalian-McKay, R., & Alterman, A. I. (1996). Gender differences in the relapse experiences of cocaine patients. *The Journal of nervous and mental disease*, 184(10), 616-622.
- McKay, B. M., Matthews, E. A., Oliveira, F. A., & Disterhoft, J. F. (2009). Intrinsic neuronal excitability is reversibly altered by a single experience in fear conditioning. *Journal of neurophysiology*, 102(5), 2763-2770.
- Milad, M. R., & Quirk, G. J. (2002). Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature*, 420(6911), 70-74.
- Milad, M. R., Igoe, S. A., Lebron-Milad, K., & Novales, J. E. (2009). Estrous cycle phase and gonadal hormones influence conditioned fear extinction. *Neuroscience*, 164(3), 887-895.
- Milad, M. R., Zeidan, M. A., Contero, A., Pitman, R. K., Klibanski, A., Rauch, S. L., & Goldstein, J. M. (2010). The influence of gonadal hormones on conditioned fear extinction in healthy humans. *Neuroscience*, 168(3), 652-658.
- Millan, E. Z., Marchant, N. J., & McNally, G. P. (2011). Extinction of drug seeking. *Behavioural brain research*, 217(2), 454-462.

- Mizuno, M., Yamada, K., Olariu, A., Nawa, H., & Nabeshima, T. (2000). Involvement of brain-derived neurotrophic factor in spatial memory formation and maintenance in a radial arm maze test in rats. *The Journal of Neuroscience*, *20*(18), 7116-7121.
- Mizuno, M., Yamada, K., He, J., Nakajima, A., & Nabeshima, T. (2003). Involvement of BDNF receptor TrkB in spatial memory formation. *Learning & memory*, *10*(2), 108-115.
- Montalbano, A., Baj, G., Papadia, D., Tongiorgi, E., & Sciancalepore, M. (2013). Blockade of BDNF signaling turns chemically-induced long-term potentiation into long-term depression. *Hippocampus*, *23*(10), 879-889.
- Monteiro, S. C., de Mattos, C. B., Ben, J., Netto, C. A., & Wyse, A. T. (2008). Ovariectomy impairs spatial memory: prevention and reversal by a soy isoflavone diet. *Metabolic brain disease*, *23*(3), 243-253.
- Moyer Jr, J. R., Thompson, L. T., & Disterhoft, J. F. (1996). Trace eyeblink conditioning increases CA1 excitability in a transient and learning-specific manner. *The Journal of Neuroscience*, *16*(17), 5536-5546.
- Moyer, J. R., Power, J. M., Thompson, L. T., & Disterhoft, J. F. (2000). Increased excitability of aged rabbit CA1 neurons after trace eyeblink conditioning. *The Journal of Neuroscience*, *20*(14), 5476-5482.
- Mozzachiodi, R., & Byrne, J. H. (2010). More than synaptic plasticity: role of nonsynaptic plasticity in learning and memory. *Trends in neurosciences*, *33*(1), 17-26.
- Mueller, D., Porter, J. T., & Quirk, G. J. (2008). Noradrenergic signaling in infralimbic cortex increases cell excitability and strengthens memory for fear extinction. *The Journal of neuroscience*, *28*(2), 369-375.
- Murphy, D. D., Cole, N. B., Greenberger, V., & Segal, M. (1998). Estradiol increases dendritic spine density by reducing GABA neurotransmission in hippocampal neurons. *The Journal of neuroscience*, *18*(7), 2550-2559.
- Myers, K. M., Carlezon, W. A., & Davis, M. (2011). Glutamate receptors in extinction and extinction-based therapies for psychiatric illness. *Neuropsychopharmacology*, *36*(1), 274-293.
- O'Neal, M. F., Means, L. W., Poole, M. C., & Hamm, R. J. (1996). Estrogen affects performance of ovariectomized rats in a two-choice water-escape working memory task. *Psychoneuroendocrinology*, *21*(1), 51-65.
- Österlund, M. K., Gustafsson, J. A., Keller, E., & Hurd, Y. L. (2000). Estrogen receptor β (ER β) messenger ribonucleic acid (mRNA) expression within the human forebrain: distinct distribution pattern to ER α mRNA 1. *The Journal of Clinical Endocrinology & Metabolism*, *85*(10), 3840-3846.

- Otis, J. M., Dashew, K. B., & Mueller, D. (2013). Neurobiological dissociation of retrieval and reconsolidation of cocaine-associated memory. *The Journal of Neuroscience*, *33*(3), 1271-1281.
- Otis, J. M., Fitzgerald, M. K., & Mueller, D. (2014). Infralimbic BDNF/TrkB enhancement of GluN2B currents facilitates extinction of a cocaine-conditioned place preference. *The Journal of Neuroscience*, *34*(17), 6057-6064.
- Packard, M. G., Kohlmaier, J. R., & Alexander, G. M. (1996). Posttraining intrahippocampal estradiol injections enhance spatial memory in male rats: interaction with cholinergic systems. *Behavioral neuroscience*, *110*(3), 626.
- Packard, M. G., & Teather, L. A. (1997). Posttraining estradiol injections enhance memory in ovariectomized rats: cholinergic blockade and synergism. *Neurobiology of learning and memory*, *68*(2), 172-188.
- Paxinos, G., & Watson, C. (2007). *The rat brain in stereotaxic coordinates* (6th ed.). Amsterdam: Elsevier.
- Pereira, L. M., Bastos, C. P., de Souza, J. M., Ribeiro, F. M., & Pereira, G. S. (2014). Estradiol enhances object recognition memory in Swiss female mice by activating hippocampal estrogen receptor α . *Neurobiology of learning and memory*, *114*, 1-9.
- Peris, J., Decambre, N., Coleman-Hardee, M. L., & Simpkins, J. W. (1991). Estradiol enhances behavioral sensitization to cocaine and amphetamine-stimulated striatal [3 H] dopamine release. *Brain research*, *566*(1), 255-264.
- Perrotti, L. I., Russo, S. J., Fletcher, H., Chin, J., Webb, T., Jenab, S., & Quinones-Jenab, V. (2001). Ovarian Hormones Modulate Cocaine-Induced Locomotor and Stereotypic Activity. *Annals of the New York Academy of Sciences*, *937*(1), 202-216.
- Perry, A. N., Westenbroek, C., & Becker, J. B. (2013). Impact of pubertal and adult estradiol treatments on cocaine self-administration. *Hormones and behavior*, *64*(4), 573-578.
- Peters, J., LaLumiere, R. T., & Kalivas, P. W. (2008). Infralimbic prefrontal cortex is responsible for inhibiting cocaine seeking in extinguished rats. *The Journal of neuroscience*, *28*(23), 6046-6053.
- Peters, J., Kalivas, P. W., & Quirk, G. J. (2009). Extinction circuits for fear and addiction overlap in prefrontal cortex. *Learning & memory*, *16*(5), 279-288.
- Peters, J., Dieppa-Perea, L. M., Melendez, L. M., & Quirk, G. J. (2010). Induction of fear extinction with hippocampal-infralimbic BDNF. *Science*, *328*(5983), 1288-1290.
- Pitha, J., & Pitha, J. (1985). Amorphous water-soluble derivatives of cyclodextrins: Nontoxic dissolution enhancing excipients. *Journal of pharmaceutical sciences*, *74*(9), 987-990.
- Pluchino, N., Ninni, F., Casarosa, E., Giannini, A., Merlini, S., Cubeddu, A., ... & Genazzani, A. R. (2009). Sex differences in brain and plasma β -endorphin content following

- testosterone, dihydrotestosterone and estradiol administration to gonadectomized rats. *Neuroendocrinology*, 89(4), 411-423.
- Poo, M. M. (2001). Neurotrophins as synaptic modulators. *Nature Reviews Neuroscience*, 2(1), 24-32.
- Quirk, G. J., Russo, G. K., Barron, J. L., & Lebron, K. (2000). The role of ventromedial prefrontal cortex in the recovery of extinguished fear. *The Journal of Neuroscience*, 20(16), 6225-6231.
- Quirk, G. J., & Mueller, D. (2008). Neural Mechanisms of Extinction Learning and Retrieval. *Neuropsychopharmacology : Official Publication of the American College of Neuropsychopharmacology*, 33(1), 56-72.
- Reichardt, L. F. (2006). Neurotrophin-regulated signalling pathways. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 361(1473), 1545-1564.
- Rhodes, M. E., & Frye, C. A. (2006). ER β -selective SERMs produce mnemonic-enhancing effects in the inhibitory avoidance and water maze tasks. *Neurobiology of learning and memory*, 85(2), 183-191.
- Rivera, R., Yacobson, I., & Grimes, D. (1999). The mechanism of action of hormonal contraceptives and intrauterine contraceptive devices. *American journal of obstetrics and gynecology*, 181(5), 1263-1269.
- Roberts, D. C. S., Bennett, S. A. L., & Vickers, G. J. (1989). The estrous cycle affects cocaine self-administration on a progressive ratio schedule in rats. *Psychopharmacology*, 98(3), 408-411.
- Robinson, T. E., & Becker, J. B. (1986). Enduring changes in brain and behavior produced by chronic amphetamine administration: a review and evaluation of animal models of amphetamine psychosis. *Brain Research Reviews*, 11(2), 157-198.
- Russo, S. J., Festa, E. D., Fabian, S. J., Gazi, F. M., Kraish, M., Jenab, S., & Quinones-Jenab, V. (2003). Gonadal hormones differentially modulate cocaine-induced conditioned place preference in male and female rats. *Neuroscience*, 120(2), 523-533.
- Russo-Neustadt, A. A., & Chen, M. J. (2005). Brain-derived neurotrophic factor and antidepressant activity. *Current pharmaceutical design*, 11(12), 1495-1510.
- Sah, P., & Davies, P. (2000). Calcium-activated potassium currents in mammalian neurons. *Clinical and Experimental Pharmacology and Physiology*, 27(9), 657-663.
- Sandstrom, N. J., & Williams, C. L. (2001). Memory retention is modulated by acute estradiol and progesterone replacement. *Behavioral neuroscience*, 115(2), 384.
- Santini, E., Muller, R. U., & Quirk, G. J. (2001). Consolidation of extinction learning involves transfer from NMDA-independent to NMDA-dependent memory. *The Journal of Neuroscience*, 21(22), 9009-9017.

- Santini, E., Ge, H., Ren, K., de Ortiz, S. P., & Quirk, G. J. (2004). Consolidation of fear extinction requires protein synthesis in the medial prefrontal cortex. *The Journal of Neuroscience*, *24*(25), 5704-5710.
- Santini, E., Quirk, G. J., & Porter, J. T. (2008). Fear conditioning and extinction differentially modify the intrinsic excitability of infralimbic neurons. *The Journal of Neuroscience*, *28*(15), 4028-4036.
- Sasahara, K., Shikimi, H., Haraguchi, S., Sakamoto, H., Honda, S. I., Harada, N., & Tsutsui, K. (2007). Mode of action and functional significance of estrogen-inducing dendritic growth, spinogenesis, and synaptogenesis in the developing Purkinje cell. *The Journal of neuroscience*, *27*(28), 7408-7417.
- Sehgal, M., Song, C., Ehlers, V. L., & Moyer, J. R. (2013). Learning to learn—intrinsic plasticity as a metaplasticity mechanism for memory formation. *Neurobiology of learning and memory*, *105*, 186-199.
- Song, C., Detert, J. A., Sehgal, M., & Moyer, J. R. (2012). Trace fear conditioning enhances synaptic and intrinsic plasticity in rat hippocampus. *Journal of neurophysiology*, *107*(12), 3397-3408.
- Sato, K., Akaishi, T., Matsuki, N., Ohno, Y., & Nakazawa, K. (2007). β -Estradiol induces synaptogenesis in the hippocampus by enhancing brain-derived neurotrophic factor release from dentate gyrus granule cells. *Brain research*, *1150*, 108-120.
- Sawai, T., Bernier, F., Fukushima, T., Hashimoto, T., Ogura, H., & Nishizawa, Y. (2002). Estrogen induces a rapid increase of calcium-calmodulin-dependent protein kinase II activity in the hippocampus. *Brain research*, *950*(1), 308-311.
- Scharfman, H. E., Mercurio, T. C., Goodman, J. H., Wilson, M. A., & MacLusky, N. J. (2003). Hippocampal excitability increases during the estrous cycle in the rat: a potential role for brain-derived neurotrophic factor. *The Journal of neuroscience*, *23*(37), 11641-11652.
- Schiess, M. C., Marian, J., & Shinnick-Gallagher, P. (1988). Estrogen priming affects active membrane properties of medial amygdala neurons. *Brain research*, *440*(2), 380-385.
- See, R. E. (2005). Neural substrates of cocaine-cue associations that trigger relapse. *European journal of pharmacology*, *526*(1), 140-146.
- Segarra, A. C., Torres-Díaz, Y. M., Silva, R. D., Puig-Ramos, A., Menéndez-Delmestre, R., Rivera-Bermúdez, J. G., ... & Agosto-Rivera, J. L. (2014). Estrogen receptors mediate estradiol's effect on sensitization and CPP to cocaine in female rats: Role of contextual cues. *Hormones and behavior*, *65*(2), 77-87.
- Shalev, U., Grimm, J. W., & Shaham, Y. (2002). Neurobiology of relapse to heroin and cocaine seeking: a review. *Pharmacological reviews*, *54*(1), 1-42.
- Sharpe, R. M. (1998). The roles of oestrogen in the male. *Trends in Endocrinology & Metabolism*, *9*(9), 371-377.

- Shughrue, P. J., Scrimo, P. J., & Merchenthaler, I. (1998). Evidence of the colocalization of estrogen receptor- β mRNA and estrogen receptor- α immunoreactivity in neurons of the rat forebrain. *Endocrinology*, *139*(12), 5267-527.
- Shughrue, P. J., & Merchenthaler, I. (2000). Evidence for novel estrogen binding sites in the rat hippocampus. *Neuroscience*, *99*(4), 605-612.
- Shughrue, P. J., & Merchenthaler, I. (2001). Distribution of estrogen receptor β immunoreactivity in the rat central nervous system. *Journal of Comparative Neurology*, *436*(1), 64-81.
- Sierra-Mercado, D., Padilla-Coreano, N., & Quirk, G. J. (2011). Dissociable roles of prelimbic and infralimbic cortices, ventral hippocampus, and basolateral amygdala in the expression and extinction of conditioned fear. *Neuropsychopharmacology*, *36*(2), 529-538.
- Singh, M., Meyer, E. M., & Simpkins, J. W. (1995). The effect of ovariectomy and estradiol replacement on brain-derived neurotrophic factor messenger ribonucleic acid expression in cortical and hippocampal brain regions of female Sprague-Dawley rats. *Endocrinology*, *136*(5), 2320-2324.
- Sircar, R., & Kim, D. (1999). Female gonadal hormones differentially modulate cocaine-induced behavioral sensitization in Fischer, Lewis, and Sprague-Dawley rats. *Journal of Pharmacology and Experimental Therapeutics*, *289*(1), 54-65.
- Sohrabji, F., Miranda, R. C., & Toran-Allerand, C. D. (1995). Identification of a putative estrogen response element in the gene encoding brain-derived neurotrophic factor. *Proceedings of the National Academy of Sciences*, *92*(24), 11110-11114.
- Solum, D. T., & Handa, R. J. (2002). Estrogen regulates the development of brain-derived neurotrophic factor mRNA and protein in the rat hippocampus. *The Journal of neuroscience*, *22*(7), 2650-2659.
- Spencer, J. L., Waters, E. M., Romeo, R. D., Wood, G. E., Milner, T. A., & McEwen, B. S. (2008). Uncovering the mechanisms of estrogen effects on hippocampal function. *Frontiers in neuroendocrinology*, *29*(2), 219-237.
- Srivastava, D. P., Waters, E. M., Mermelstein, P. G., Kramár, E. A., Shors, T. J., & Liu, F. (2011). Rapid estrogen signaling in the brain: implications for the fine-tuning of neuronal circuitry. *The Journal of Neuroscience*, *31*(45), 16056-16063.
- Srivastava, D. P., & Evans, P. D. (2013). G-Protein Oestrogen Receptor 1: Trials and Tribulations of a Membrane Oestrogen Receptor. *Journal of neuroendocrinology*, *25*(11), 1219-1230.
- Srivastava, D. P., Woolfrey, K. M., & Penzes, P. (2013). Insights into rapid modulation of neuroplasticity by brain estrogens. *Pharmacological reviews*, *65*(4), 1318-1350.
- Storm, J. F. (1990). Potassium currents in hippocampal pyramidal cells. *Progress in brain research*, *83*, 161-187.

- Strakowski, S. M., Sax, K. W., Setters, M. J., & Keck, P. E. (1996). Enhanced response to repeated d-amphetamine challenge: evidence for behavioral sensitization in humans. *Biological psychiatry*, *40*(9), 872-880.
- Substance Abuse and Mental Health Services Administration. Results from the 2015. National Survey on Drug Use and Health: National Findings. Rockville: 2007. (Office of Applied Studies, NSDUH Series H-30, DHHS Publication No. SMA 06-4194). MD. 2015.
- Teyler, T. J., Vardaris, R. M., Lewis, D., & Rawitch, A. B. (1980). Gonadal steroids: effects on excitability of hippocampal pyramidal cells. *Science*, *209*(4460), 1017-1018.
- Thibault, O., & Landfield, P. W. (1996). Increase in single L-type calcium channels in hippocampal neurons during aging. *Science*, *272*(5264), 1017.
- Thompson, L. T., Moyer, J. R., & Disterhoft, J. F. (1996). Transient changes in excitability of rabbit CA3 neurons with a time course appropriate to support memory consolidation. *Journal of Neurophysiology*, *76*(3), 1836-1849.
- Tombaugh, G. C., Rowe, W. B., & Rose, G. M. (2005). The slow afterhyperpolarization in hippocampal CA1 neurons covaries with spatial learning ability in aged Fisher 344 rats. *The Journal of neuroscience*, *25*(10), 2609-2616.
- Toran-Allerand, C. D., Singh, M., & Sétáló, G. (1999). Novel mechanisms of estrogen action in the brain: new players in an old story. *Frontiers in neuroendocrinology*, *20*(2), 97-121.
- Torregrossa, M. M., & Taylor, J. R. (2013). Learning to forget: manipulating extinction and reconsolidation processes to treat addiction. *Psychopharmacology*, *226*(4), 659-672.
- Tuscher, J. J., Fortress, A. M., Kim, J., & Frick, K. M. (2015). Regulation of object recognition and object placement by ovarian sex steroid hormones. *Behavioural brain research*, *285*, 140-157.
- Tuscher, J. J., Luine, V., Frankfurt, M., & Frick, K. M. (2016). Estradiol-mediated spine changes in the dorsal hippocampus and medial prefrontal cortex of ovariectomized female mice depend on ERK and mTOR activation in the dorsal hippocampus. *The Journal of Neuroscience*, *36*(5), 1483-1489.
- Twining, R. C., Tuscher, J. J., Doncheck, E. M., Frick, K. M., & Mueller, D. (2013). 17 β -Estradiol is necessary for extinction of cocaine seeking in female rats. *Learning & memory*, *20*(6), 300-306.
- Tzschentke, T. M. (1998). Measuring reward with the conditioned place preference paradigm: a comprehensive review of drug effects, recent progress and new issues. *Progress in neurobiology*, *56*(6), 613-672.
- Tzschentke, T. M. (2007). Review on CPP: Measuring reward with the conditioned place preference (CPP) paradigm: update of the last decade. *Addiction biology*, *12*(3-4), 227-462.

- Vassoler, F. M., White, S. L., Schmidt, H. D., Sadri-Vakili, G., & Pierce, R. C. (2013). Epigenetic inheritance of a cocaine-resistance phenotype. *Nature neuroscience*, *16*(1), 42-47.
- Vasudevan, N., & Pfaff, D. W. (2007). Membrane-initiated actions of estrogens in neuroendocrinology: emerging principles. *Endocrine reviews*, *28*(1), 1-19.
- Walf, A. A., Rhodes, M. E., & Frye, C. A. (2006). Ovarian steroids enhance object recognition in naturally cycling and ovariectomized, hormone-primed rats. *Neurobiology of learning and memory*, *86*(1), 35-46.
- Walf, A. A., Koonce, C. J., & Frye, C. A. (2008). Estradiol or diarylpropionitrile administration to wild type, but not estrogen receptor beta knockout, mice enhances performance in the object recognition and object placement tasks. *Neurobiology of learning and memory*, *89*(4), 513-521.
- Wallace, M., Luine, V., Arellanos, A., & Frankfurt, M. (2006). Ovariectomized rats show decreased recognition memory and spine density in the hippocampus and prefrontal cortex. *Brain research*, *1126*(1), 176-182.
- Weiss, R. D., Martinez-Raga, J., Griffin, M. L., Greenfield, S. F., & Hufford, C. (1997). Gender differences in cocaine dependent patients: a 6 month follow-up study. *Drug and Alcohol Dependence*, *44*(1), 35-40.
- Westermeyer, J., & Boedicker, A. E. (2000). Course, severity, and treatment of substance abuse among women versus men. *The American journal of drug and alcohol abuse*, *26*(4), 523-535.
- Wong, M., & Moss, R. L. (1991). Electrophysiological evidence for a rapid membrane action of the gonadal steroid, 17 β -estradiol, on CA1 pyramidal neurons of the rat hippocampus. *Brain research*, *543*(1), 148-152.
- Wong, M., & Moss, R. L. (1992). Long-term and short-term electrophysiological effects of estrogen on the synaptic properties of hippocampal CA1 neurons. *The Journal of neuroscience*, *12*(8), 3217-3225.
- Woolley, C. S., & McEwen, B. S. (1993). Roles of estradiol and progesterone in regulation of hippocampal dendritic spine density during the estrous cycle in the rat. *Journal of Comparative Neurology*, *336*(2), 293-306.
- Woolley, C. S. (1998). Estrogen-mediated structural and functional synaptic plasticity in the female rat hippocampus. *Hormones and behavior*, *34*(2), 140-148.
- Woolley, C. S. (2007). Acute effects of estrogen on neuronal physiology. *Annu. Rev. Pharmacol. Toxicol.*, *47*, 657-680.
- Wouda, J. A., Diergaarde, L., Riga, D., Van Mourik, Y., Schoffelmeer, A. N., & De Vries, T. J. (2010). Disruption of long-term alcohol-related memory reconsolidation: role of β -adrenoceptors and NMDA receptors. *Frontiers in behavioral neuroscience*, *4*, 179.

- Wu, W. W., Adelman, J. P., & Maylie, J. (2011). Ovarian hormone deficiency reduces intrinsic excitability and abolishes acute estrogen sensitivity in hippocampal CA1 pyramidal neurons. *The Journal of Neuroscience*, *31*(7), 2638-2648.
- Wu, Y. C., Hill, R. A., Gogos, A., & Van Den Buuse, M. (2013). Sex differences and the role of estrogen in animal models of schizophrenia: interaction with BDNF. *Neuroscience*, *239*, 67-83.
- Xin, J., Ma, L., Zhang, T. Y., Yu, H., Wang, Y., Kong, L., & Chen, Z. Y. (2014). Involvement of BDNF signaling transmission from basolateral amygdala to infralimbic prefrontal cortex in conditioned taste aversion extinction. *The Journal of Neuroscience*, *34*(21), 7302-7313.
- Yamada, K., & Nabeshima, T. (2003). Brain-derived neurotrophic factor/TrkB signaling in memory processes. *Journal of pharmacological sciences*, *91*(4), 267-270.
- Ying, S. W., Futter, M., Rosenblum, K., Webber, M. J., Hunt, S. P., Bliss, T. V., & Bramham, C. R. (2002). Brain-derived neurotrophic factor induces long-term potentiation in intact adult hippocampus: requirement for ERK activation coupled to CREB and upregulation of Arc synthesis. *The Journal of neuroscience*, *22*(5), 1532-1540.
- Yoshii, A., & Constantine-Paton, M. (2010). Postsynaptic BDNF-TrkB signaling in synapse maturation, plasticity, and disease. *Developmental neurobiology*, *70*(5), 304-322.
- Zeidan, M. A., Igoe, S. A., Linnman, C., Vitalo, A., Levine, J. B., Klibanski, A., ... & Milad, M. R. (2011). Estradiol modulates medial prefrontal cortex and amygdala activity during fear extinction in women and female rats. *Biological psychiatry*, *70*(10), 920-927.
- Zhao, Z., Fan, L., & Frick, K. M. (2010). Epigenetic alterations regulate estradiol-induced enhancement of memory consolidation. *Proceedings of the National Academy of Sciences*, *107*(12), 5605-5610.
- Zhang, W., & Linden, D. J. (2003). The other side of the engram: experience-driven changes in neuronal intrinsic excitability. *Nature Reviews Neuroscience*, *4*(11), 885-900.
- Zhang, J. C., Yao, W., Dong, C., Yang, C., Ren, Q., Ma, M., ... & Hashimoto, K. (2015). Comparison of ketamine, 7, 8-dihydroxyflavone, and ANA-12 antidepressant effects in the social defeat stress model of depression. *Psychopharmacology*, *232*(23), 4325-4335.
- Zhou, J., Zhang, H., Cohen, R. S., & Pandey, S. C. (2005). Effects of estrogen treatment on expression of brain-derived neurotrophic factor and cAMP response element-binding protein expression and phosphorylation in rat amygdaloid and hippocampal structures. *Neuroendocrinology*, *81*(5), 294-310.
- Znamensky, V., Akama, K. T., McEwen, B. S., & Milner, T. A. (2003). Estrogen levels regulate the subcellular distribution of phosphorylated Akt in hippocampal CA1 dendrites. *The Journal of neuroscience*, *23*(6), 2340-2347.